



ELSEVIER

Contents lists available at ScienceDirect

Journal of Functional Analysis

www.elsevier.com/locate/jfa



Scattering matrices and Dirichlet-to-Neumann maps



Jussi Behrndt^{a,*}, Mark M. Malamud^{b,c}, Hagen Neidhardt^d

^a *Institut für Numerische Mathematik, Technische Universität Graz, Steyrergasse 30, 8010 Graz, Austria*

^b *Institute of Applied Mathematics and Mechanics, National Academy of Science of Ukraine, Dobrovol's'kogo Str. 1, 84100 Slavyansk, Donetsk Region, Ukraine*

^c *Peoples Friendship University of Russia (RUDN University), 6 Miklukho-Maklaya Street, Moscow, 117198, Russian Federation*

^d *Weierstrass Institute for Applied Analysis and Stochastics, Mohrenstr. 39, 10117 Berlin, Germany*

ARTICLE INFO

Article history:

Received 23 July 2016

Accepted 1 June 2017

Available online 10 June 2017

Communicated by B. Schlein

MSC:

35J10

35P25

47A40

81U20

Keywords:

Scattering matrix

Titchmarsh–Weyl m -function

Schrödinger operator

Dirichlet-to-Neumann map

ABSTRACT

A general representation formula for the scattering matrix of a scattering system consisting of two self-adjoint operators in terms of an abstract operator valued Titchmarsh–Weyl m -function is proved. This result is applied to scattering problems for different self-adjoint realizations of Schrödinger operators on unbounded domains, Schrödinger operators with singular potentials supported on hypersurfaces, and orthogonal couplings of Schrödinger operators. In these applications the scattering matrix is expressed in an explicit form with the help of Dirichlet-to-Neumann maps.

© 2017 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail addresses: behrndt@tugraz.at (J. Behrndt), malamud3m@gmail.com (M.M. Malamud), neidhard@wias-berlin.de (H. Neidhardt).

1. Introduction

Let A and B be self-adjoint operators in a Hilbert space \mathfrak{H} and assume that the resolvent difference

$$(B - \lambda)^{-1} - (A - \lambda)^{-1}, \quad \lambda \in \rho(A) \cap \rho(B), \quad (1.1)$$

belongs to the ideal $\mathfrak{S}_1(\mathfrak{H})$ of trace class operators. It is well known that in this situation the wave operators $W_{\pm}(A, B)$ of the pair $\{A, B\}$ exist and are complete, and the scattering operator $S(A, B) = W_+(A, B)^* W_-(A, B)$ is unitarily equivalent to a multiplication operator induced by a family $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ of unitary operators $S(A, B; \lambda)$ in the spectral representation of the absolutely continuous part of A . This family is called the scattering matrix of the scattering system $\{A, B\}$ and is one of the most important quantities in the analysis of scattering processes; we refer the reader to the monographs [12, 59, 79, 81, 82] for more details.

The main objective of this paper is to express the scattering matrix of $\{A, B\}$ in terms of an abstract operator valued Titchmarsh–Weyl m -function, and to apply this result to scattering problems for Schrödinger operators. In order to explain our main abstract result [Theorem 3.1](#) consider the closed symmetric operator $S = A \cap B$ and note that S has infinite defect numbers whenever the resolvent difference of A and B in (1.1) is infinite dimensional. The closure of the operator $T = A \hat{+} B$, where $\hat{+}$ denotes the sum of subspaces in $\mathfrak{H} \times \mathfrak{H}$, coincides with S^* and clearly A and B are self-adjoint restrictions of T . This setting can be fitted in the framework of (B) -generalized boundary triples or quasi boundary triples and their Weyl functions from [38] or [13, 14], respectively, and allows to introduce boundary maps Γ_0 and Γ_1 on $\text{dom}(T)$, which can be viewed as abstract analogs of the Dirichlet and Neumann trace operators (see also [34, 35]). For $\lambda \in \mathbb{C} \setminus \mathbb{R}$ one defines the Weyl function M via

$$M(\lambda)\Gamma_0 f_\lambda = \Gamma_1 f_\lambda, \quad f_\lambda \in \ker(T - \lambda),$$

see [Section 2](#) for the details. In PDE applications $M(\lambda)$ is usually the Dirichlet-to-Neumann map (or its inverse, the Neumann-to-Dirichlet map) acting in some boundary space. Roughly speaking our main abstract result states that the scattering matrix of $\{A, B\}$ is of the form

$$S(A, B; \lambda) = I - 2i\sqrt{\text{Im } M(\lambda + i0)} M(\lambda + i0)^{-1} \sqrt{\text{Im } M(\lambda + i0)}$$

for a.e. $\lambda \in \mathbb{R}$. This representation is a highly nontrivial generalization of a similar result from [19], where the special case that the resolvent difference in (1.1) is a finite rank operator was treated in the context of ordinary boundary triples and their Weyl functions from [37, 38], see also [2], [8, Chapter 4], [82, Chapter 3, §1], and [20] for related results and simple examples. In contrast to the earlier results in the finite rank case the present representation formula is applicable to scattering problems for Schrödinger operators (or

more general elliptic second order differential operators) on unbounded domains, which we shall explain in more detail next.

In fact, our main motivation for establishing the general representation formula for the scattering matrix in Section 3 in an abstract extension theory framework is the applicability to scattering problems for Schrödinger operators with Dirichlet, Neumann, and Robin boundary conditions on exterior domains in \mathbb{R}^2 and \mathbb{R}^3 in Section 4, and orthogonal couplings of Schrödinger operators, and Schrödinger operators with singular potentials supported on curves and hypersurfaces in \mathbb{R}^2 and \mathbb{R}^3 in Section 5. Let us first explain the situation for a scattering system consisting of a Neumann and a Robin realization; for more details and a slightly more general situation see Section 4.4. Denote the Dirichlet and Neumann trace operators by γ_D and γ_N , respectively, and consider the self-adjoint operators

$$Af = -\Delta f + Vf, \quad \text{dom}(A) = \{f \in H^2(\Omega) : \gamma_N f = 0\},$$

and

$$Bf = -\Delta f + Vf, \quad \text{dom}(B) = \{f \in H^2(\Omega) : \alpha \gamma_D f = \gamma_N f\},$$

where $\alpha \in C^2(\partial\Omega)$ is real, the potential V is real and bounded, and the domain Ω is the complement of a bounded set with a C^∞ -smooth boundary in \mathbb{R}^2 or \mathbb{R}^3 . In this situation it is known from [15,58] that the resolvent difference of A and B satisfies the trace class condition (1.1). If $\mathcal{N}(\lambda)$, $\lambda \in \mathbb{C} \setminus \mathbb{R}$, denotes the Neumann-to-Dirichlet map, that is,

$$\mathcal{N}(\lambda)\gamma_N f_\lambda = \gamma_D f_\lambda, \quad -\Delta f_\lambda + Vf_\lambda = \lambda f_\lambda,$$

we obtain in Theorem 4.7 that the scattering matrix of the scattering system $\{A, B\}$ admits the form

$$S(A, B; \lambda) = I_{\mathcal{G}_\lambda} + 2i\sqrt{\text{Im}\mathcal{N}(\lambda + i0)}(I - \alpha\mathcal{N}(\lambda + i0))^{-1}\alpha\sqrt{\text{Im}\mathcal{N}(\lambda + i0)}$$

for a.e. $\lambda \in \mathbb{R}$. Here the space $L^2(\mathbb{R}, d\lambda, \mathcal{G}_\lambda)$, where $\mathcal{G}_\lambda = \overline{\text{ran}(\text{Im}\mathcal{N}(\lambda + i0))}$ for a.e. $\lambda \in \mathbb{R}$, forms a spectral representation of the absolutely continuous part of the Neumann operator A_N and the limits $\text{Im}\mathcal{N}(\lambda + i0)$ and $(I - \alpha\mathcal{N}(\lambda + i0))^{-1}$ have to be interpreted in suitable operator topologies; cf. Theorem 4.7 for details. A similar result is proved in Theorem 4.3 for the pair consisting of the Dirichlet realization of $-\Delta + V$ and the Robin operator B in $L^2(\mathbb{R}^2)$; here the trace class property (1.1) for $n = 2$ is due to Birman [24]. For some recent work on related spectral problems for Schrödinger operators we refer the reader to [9,22,30,46–50,64,67,74,77] and for more general partial elliptic differential operators to [1,13,14,17,18,21,29,55–58,63,65,66,75,76].

Our second set of examples in Section 5 is a bit more involved. Here scattering systems consisting of the free Schrödinger operator

$$Af = -\Delta f + Vf, \quad \text{dom}(A) = H^2(\mathbb{R}^n), \quad (1.2)$$

and orthogonal couplings of Schrödinger operators with Dirichlet and Neumann boundary conditions, or Schrödinger operators with singular δ -potentials of strength $\alpha \in L^\infty(\mathcal{C})$ supported on hypersurfaces \mathcal{C} which split \mathbb{R}^2 or \mathbb{R}^3 into a bounded smooth domain Ω_+ and a smooth exterior domain Ω_- are studied. The latter operator is of the form

$$Bf = -\Delta f + Vf, \quad \text{dom}(B) = \left\{ f = \begin{pmatrix} f_+ \\ f_- \end{pmatrix} \in H_{\Delta}^{3/2}(\mathbb{R}^n \setminus \mathcal{C}) : \begin{array}{l} \gamma_D^+ f_+ = \gamma_D^- f_-, \\ \alpha \gamma_D^\pm f_\pm = \gamma_N^+ f_+ + \gamma_N^- f_- \end{array} \right\}; \quad (1.3)$$

here $H_{\Delta}^{3/2}(\mathbb{R}^n \setminus \mathcal{C})$ is a subspace of $H^{3/2}(\Omega_+) \times H^{3/2}(\Omega_-)$ and γ_D^\pm and γ_N^\pm denote the Dirichlet and Neumann trace operators on the interior and exterior domain; cf. Section 5.4 for the details. Schrödinger operators with δ -potentials play an important role in various physically relevant problems and have therefore attracted a lot of attention. We refer the interested reader to the review paper [39], to e.g. [7,10,16,27,40–43] and the monographs [6,8] for more details and further references. We shall briefly discuss the scattering matrix for the pair of operators in (1.2)–(1.3); for the pairs consisting of A in (1.2) and the orthogonal sum of the Dirichlet or the Neumann realizations of $-\Delta + V$ on Ω_+ and Ω_- see Theorem 5.1 and Theorem 5.4, respectively. It follows from [16] that the above choice of A and B satisfies the trace class condition (1.1) in dimensions $n = 2$ and $n = 3$ and we show in this situation in Theorem 5.6 that the scattering matrix is given by

$$S(A, B; \lambda) = I_{\mathcal{G}_\lambda} + 2i\sqrt{\text{Im } \mathcal{E}(\lambda + i0)}(I - \alpha \mathcal{E}(\lambda + i0))^{-1} \alpha \sqrt{\text{Im } \mathcal{E}(\lambda + i0)},$$

where the function \mathcal{E} is defined as

$$\mathcal{E}(\lambda) = (\mathcal{D}_+(\lambda)^{-1} + \mathcal{D}_-(\lambda)^{-1})^{-1}, \quad \lambda \in \mathbb{C} \setminus \mathbb{R},$$

and $\mathcal{D}_\pm(\lambda)$ denote the Dirichlet-to-Neumann maps corresponding to $-\Delta + V$ on the domains Ω_\pm . In this context we also refer the reader to related work by Pavlov and coauthors in [11,69,72], where scattering problems for certain couplings of Schrödinger operators were considered.

1.1. Notation

Throughout the paper \mathfrak{H} and \mathcal{H} denote separable Hilbert spaces with scalar product (\cdot, \cdot) . The linear space of bounded linear operators defined from \mathfrak{H} to \mathcal{H} is denoted by $\mathcal{B}(\mathfrak{H}, \mathcal{H})$. For brevity we write $\mathcal{B}(\mathfrak{H})$ instead of $\mathcal{B}(\mathfrak{H}, \mathfrak{H})$. The ideal of compact operators is denoted by $\mathfrak{S}_\infty(\mathfrak{H}, \mathcal{H})$ and $\mathfrak{S}_\infty(\mathfrak{H})$. For $p > 0$ the Schatten–von Neumann ideals are denoted by $\mathfrak{S}_p(\mathfrak{H}, \mathcal{H})$ and $\mathfrak{S}_p(\mathfrak{H})$; they consist of all compact operators T with

p -summable singular values $s_j(T)$ (i.e. eigenvalues of $(T^*T)^{1/2}$). We shall also work with the operator ideals

$$\mathcal{S}_p(\mathfrak{H}, \mathcal{H}) = \{T \in \mathfrak{S}_\infty(\mathfrak{H}, \mathcal{H}) \mid s_j(T) = O(j^{-1/p}) \text{ as } j \rightarrow \infty\}, \quad p > 0,$$

and we recall that

$$\mathcal{S}_p(\mathcal{H}, \mathfrak{H}) \cdot \mathcal{S}_q(\mathfrak{H}, \mathcal{H}) = \mathcal{S}_r(\mathfrak{H}), \quad \text{where} \quad \frac{1}{p} + \frac{1}{q} = \frac{1}{r}. \quad (1.4)$$

The resolvent set and the spectrum of a linear operator A is denoted by $\rho(A)$ and $\sigma(A)$, respectively. The domain, kernel and range of a linear operator A are denoted by $\text{dom}(A)$, $\ker(A)$, and $\text{ran}(A)$, respectively. By $\mathfrak{B}(\mathbb{R})$ we denote the Borel sets of \mathbb{R} . The Lebesgue measure on $\mathfrak{B}(\mathbb{R})$ is denoted by $d\lambda$.

A holomorphic function $M(\cdot) : \mathbb{C}_+ \rightarrow \mathcal{B}(\mathcal{H})$ is a Nevanlinna (or Herglotz or R -function) if its imaginary part $\text{Im}(M(z)) := \frac{1}{2i}(M(z) - M(z)^*)$, $z \in \mathbb{C}_+$, is a non-negative operator. Nevanlinna functions are extended to \mathbb{C}_- by $M(z) := M(\bar{z})^*$, $z \in \mathbb{C}_-$. The class of $\mathcal{B}(\mathcal{H})$ -valued Nevanlinna functions is denoted by $R[\mathcal{H}]$. A Nevanlinna function satisfying $\ker(\text{Im}(M(z))) = \{0\}$ ($0 \in \rho(\text{Im}(M(z)))$) for some, and hence for all, $z \in \mathbb{C}_+$, is said to be strict (uniformly strict, respectively). These subclasses are denoted by $R^s[\mathcal{H}]$ and $R^u[\mathcal{H}]$, respectively.

2. Self-adjoint extensions of symmetric operators and abstract Titchmarsh–Weyl m -functions

In the preparatory Section 2.1 we recall the notion of boundary triples and their Weyl functions from extension theory of symmetric operators, and we introduce the concept of \mathfrak{S}_p -regular Weyl functions in Section 2.2. This notion is important and useful for our purposes since it is directly related (and in some situations equivalent) to the \mathfrak{S}_p -property of the resolvent difference of certain self-adjoint extensions.

2.1. B -generalized boundary triples and their Weyl functions

In this subsection we review the notion of generalized (or B -generalized) and ordinary boundary triples from extension theory of symmetric operators, and we introduce a new concept, the so-called double B -generalized boundary triples in Definition 2.1 below. We refer the reader to [28,31,34,37,38,51,80] for more details on ordinary and B -generalized boundary triples, see also [13,14,32] for related notions.

In the following S denotes a densely defined, closed, symmetric operator in a separable Hilbert space \mathfrak{H} .

Definition 2.1 ([38]). A triple $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is called a B -generalized boundary triple for S^* if \mathcal{H} is a Hilbert space and for some operator T in \mathfrak{H} such that $\overline{T} = S^*$, the linear mappings $\Gamma_0, \Gamma_1 : \text{dom}(T) \rightarrow \mathcal{H}$ satisfy the abstract Green's identity

$$(Tf, g) - (f, Tg) = (\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g), \quad f, g \in \operatorname{dom}(T), \quad (2.1)$$

the operator $A_0 := T \upharpoonright \ker(\Gamma_0)$ is self-adjoint in \mathfrak{H} , and $\operatorname{ran}(\Gamma_0) = \mathcal{H}$ holds.

If, in addition, the operator $A_1 := T \upharpoonright \ker(\Gamma_1)$ is self-adjoint in \mathfrak{H} and $\operatorname{ran}(\Gamma_1) = \mathcal{H}$, then the triple $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is called a *double B -generalized boundary triple* for S^* .

We note that a B -generalized boundary triple for S^* exists if and only if S admits self-adjoint extensions in \mathfrak{H} , that is, the deficiency indices of S coincide. Furthermore, if $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is a B -generalized boundary triple for S^* then

$$\operatorname{dom}(S) = \ker(\Gamma_0) \cap \ker(\Gamma_1)$$

holds, the mappings $\Gamma_0, \Gamma_1 : \operatorname{dom}(T) \rightarrow \mathcal{H}$ are closable when viewed as linear operators from $\operatorname{dom} S^*$ equipped with the graph norm to \mathcal{H} , and $\operatorname{ran}(\Gamma_1)$ turns out to be dense in \mathcal{H} ; cf. [38, Section 6].

The notion of double B -generalized boundary triples is inspired by the fact that the mappings in the so-called transposed triple $\Pi^\top := \{\mathcal{H}, \Gamma_1, -\Gamma_0\}$ satisfy the abstract Green's identity but since in general neither $A_1 = T \upharpoonright \ker(\Gamma_1)$ is self-adjoint nor $\operatorname{ran}(\Gamma_1) = \mathcal{H}$ holds the transposed triple Π^\top is not a B -generalized boundary triple in general. In fact, a B -generalized boundary triple $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ for S^* is a double B -generalized boundary triple for S^* if and only if the transposed triple $\Pi^\top = \{\mathcal{H}, \Gamma_1, -\Gamma_0\}$ is also a B -generalized boundary triple for S^* .

In some of the proofs of the results in Section 2.2 we shall also make use of the notion of ordinary boundary triples, which we recall here for the convenience of the reader.

Definition 2.2. A triple $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is called an *ordinary boundary triple* for S^* if \mathcal{H} is a Hilbert space, the linear mappings $\Gamma_0, \Gamma_1 : \operatorname{dom}(S^*) \rightarrow \mathcal{H}$ satisfy the abstract Green's identity

$$(S^* f, g) - (f, S^* g) = (\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g), \quad f, g \in \operatorname{dom}(S^*), \quad (2.2)$$

and the mapping $\Gamma = (\Gamma_0, \Gamma_1)^\top : \operatorname{dom}(S^*) \rightarrow \mathcal{H} \times \mathcal{H}$ is surjective.

Observe that any ordinary boundary triple is automatically a double B -generalized boundary triple; the converse is not true in general. Ordinary boundary triples are an efficient tool in extension theory of symmetric operators. In particular, if $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is an ordinary boundary triple for S^* , then all closed proper extensions $\tilde{S} \subset S^*$ of S in \mathfrak{H} can be parametrized by means of the set of closed linear relations in \mathcal{H} via

$$\tilde{S} \mapsto \Theta := \{ \{ \Gamma_0 f, \Gamma_1 f \} : f \in \operatorname{dom}(\tilde{S}) \} \subset \mathcal{H} \times \mathcal{H}. \quad (2.3)$$

We write $\tilde{S} = S_\Theta$. If Θ is an operator then (2.3) takes the form

$$S_\Theta = S^* \upharpoonright \ker(\Gamma_1 - \Theta \Gamma_0)$$

One verifies $(S_\Theta)^* = S_{\Theta^*}$ and hence the self-adjoint extensions of S in \mathfrak{H} correspond to the self-adjoint relations Θ in \mathcal{H} . We shall use that Θ in (2.3) is an operator (and not a multivalued linear relation) if and only if the extension S_Θ and $A_0 = S^* \upharpoonright \ker(\Gamma_0)$ are disjoint, that is, $A_0 \cap S_\Theta = S$.

Next we recall the notions and some important properties of γ -fields and Weyl functions. For an ordinary boundary triple they go back to [36,37], for B -generalized boundary triples we refer the reader to [38]. In the following let $\{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for S^* ; the special case of an ordinary boundary triple is then covered as well. Observe first that for each $z \in \rho(A_0)$, $A_0 = T \upharpoonright \ker(\Gamma_0)$, the following direct sum decomposition holds

$$\operatorname{dom}(T) = \operatorname{dom}(A_0) \dot{+} \ker(T - z) = \ker(\Gamma_0) \dot{+} \ker(T - z). \quad (2.4)$$

Hence the restriction of the mapping Γ_0 to $\ker(T - z)$ is injective for all $z \in \rho(A_0)$.

Definition 2.3 ([38]). Let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple. The γ -field $\gamma(\cdot)$ and Weyl function $M(\cdot)$ corresponding to Π are defined by

$$\gamma(z) := (\Gamma_0 \upharpoonright \ker(T - z))^{-1} \quad \text{and} \quad M(z) := \Gamma_1 \gamma(z), \quad z \in \rho(A_0),$$

respectively.

It follows from (2.4) that for $z \in \rho(A_0)$ the values $\gamma(z)$ of the γ -field and the values $M(z)$ of the Weyl function are both well defined linear operators on $\operatorname{ran}(\Gamma_0) = \mathcal{H}$. Moreover, $\gamma(z) \in \mathcal{B}(\mathcal{H}, \mathfrak{H})$ maps onto $\ker(T - z) \subset \ker(S^* - z) \subset \mathfrak{H}$ and for all $z, \xi \in \rho(A_0)$ the relations

$$\gamma(z) = (I + (z - \xi)(A_0 - z)^{-1})\gamma(\xi) = (A_0 - \xi)(A_0 - z)^{-1}\gamma(\xi) \quad (2.5)$$

and

$$\gamma(z)^* = \Gamma_1(A_0 - \bar{z})^{-1} \in \mathcal{B}(\mathfrak{H}, \mathcal{H}) \quad (2.6)$$

hold. In particular, $\operatorname{ran}(\gamma(z)^*) = \operatorname{ran}(\Gamma_1 \upharpoonright \operatorname{dom}(A_0))$ does not depend on the point $z \in \rho(A_0)$ and

$$(\operatorname{ran} \gamma(z)^*)^\perp = \ker \gamma(z) = \{0\}$$

shows that $\operatorname{ran}(\gamma(z)^*)$ is dense in \mathcal{H} for all $z \in \rho(A_0)$. Furthermore, it follows from (2.5) that $\gamma(\cdot)$ is holomorphic on $\rho(A_0)$.

The values of the Weyl function $M(\cdot)$ are operators in $\mathcal{B}(\mathcal{H})$ and $M(z)$ maps \mathcal{H} into the dense subspace $\operatorname{ran}(\Gamma_1) \subset \mathcal{H}$. The Weyl function and γ -field are related by the identity

$$M(z) - M(\xi)^* = (z - \bar{\xi})\gamma(\xi)^*\gamma(z), \quad z, \xi \in \rho(A_0), \quad (2.7)$$

and, in particular, $M(\bar{z}) = M(z)^*$ for all $z \in \rho(A_0)$. It follows from (2.5) and (2.7) that $M(\cdot)$ is holomorphic on $\rho(A_0)$. Setting $\xi = z$ in (2.7) one gets

$$\operatorname{Im} M(z) = \frac{1}{2i}(M(z) - M(z)^*) = (\operatorname{Im} z) \gamma(z)^* \gamma(z) \quad (2.8)$$

and hence $\operatorname{Im} M(z) \geq 0$ for $z \in \mathbb{C}_+$. This identity also yields

$$\ker(\operatorname{Im} M(z)) = \ker(\gamma(z)) = \{0\}, \quad z \in \mathbb{C}_\pm,$$

and together with the holomorphy of $M(\cdot)$ on $\rho(A_0)$ we conclude that $M(\cdot)$ is a so-called *strict* Nevanlinna function with values in $\mathcal{B}(\mathcal{H})$ (in symbols $M(\cdot) \in R^s[\mathcal{H}]$). If Π is a double B -generalized boundary triple then the Weyl function corresponding to the transposed B -generalized boundary triple $\Pi^\top = \{\mathcal{H}, \Gamma_1, -\Gamma_0\}$ is given by $-M(\cdot)^{-1}$ and also belongs to the class $R^s[\mathcal{H}]$, in particular, for $z \in \rho(A_0) \cap \rho(A_1)$ the values $M(z)$ of the Weyl function of a double B -generalized boundary triple are bounded and boundedly invertible operators.

If Π is an ordinary boundary triple then the operators $\gamma(z)$ are boundedly invertible when viewed as operators from \mathcal{H} onto $\ker(S^* - z)$. In this case it follows from (2.8) that $\operatorname{Im} M(z)$ is a uniformly positive operator for $z \in \mathbb{C}_+$, and hence the Weyl function corresponding to an ordinary boundary triple belongs to the class $R^u[\mathcal{H}]$ of the so-called *uniformly strict* Nevanlinna functions with values in $\mathcal{B}(\mathcal{H})$; cf. [34].

2.2. Resolvent comparability and \mathfrak{S}_p -regular Weyl functions

Let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for S^* with the corresponding Weyl function $M(\cdot)$, and let $A_0 = S^* \upharpoonright \ker(\Gamma_0)$ and $A_1 = S^* \upharpoonright \ker(\Gamma_1)$. It is important to characterize the property of the resolvent comparability of the operators A_0 and A_1 in terms of the Weyl function $M(\cdot)$. To this end we introduce the notion of \mathfrak{S}_p -regular Nevanlinna functions in the next definition.

Definition 2.4. A Nevanlinna function $M(\cdot) \in R[\mathcal{H}]$ is called \mathfrak{S}_p -regular for some $p \in (0, \infty]$ if it admits a representation

$$M(z) = C + K(z), \quad K(\cdot) : \mathbb{C}_+ \longrightarrow \mathfrak{S}_p(\mathcal{H}), \quad z \in \mathbb{C}_+, \quad (2.9)$$

where $C \in \mathcal{B}(\mathcal{H})$ is a self-adjoint operator such that $0 \in \rho(C)$ and $K(\cdot)$ is a strict Nevanlinna function with values in $\mathcal{B}(\mathcal{H})$, that is, $K(\cdot) \in R^s[\mathcal{H}]$. The class of \mathfrak{S}_p -regular Nevanlinna functions is denoted by $R_{\mathfrak{S}_p}^{\operatorname{reg}}[\mathcal{H}]$.

In other words, a Nevanlinna function is \mathfrak{S}_p -regular if it differs from a strict Nevanlinna function with values in \mathfrak{S}_p by a bounded and boundedly invertible self-adjoint constant.

Lemma 2.5. If $M(\cdot) \in R_{\mathfrak{S}_p}^{\operatorname{reg}}[\mathcal{H}]$ for some $p \in (0, \infty]$, then $-M(\cdot)^{-1} \in R_{\mathfrak{S}_p}^{\operatorname{reg}}[\mathcal{H}]$.

Proof. Since $M(\cdot) \in R_{\mathfrak{S}_p}^{\text{reg}}[\mathcal{H}]$ for some $p \in (0, \infty]$, there exists a boundedly invertible self-adjoint operator C and a strict Nevanlinna function $K(\cdot) \in R^s[\mathcal{H}]$ such that

$$M(z) = C + K(z), \quad z \in \mathbb{C}_+. \quad (2.10)$$

Observe first that $\ker(M(z)) = \{0\}$ holds for all $z \in \mathbb{C}_+$. In fact, $M(z)\varphi = 0$ yields $((C + \operatorname{Re} K(z))\varphi, \varphi) = 0$ and $(\operatorname{Im} K(z)\varphi, \varphi) = 0$, and as $K(\cdot)$ is strict we conclude $\varphi = 0$ from the latter. Furthermore, as $0 \in \rho(C)$ and $K(z) \in \mathfrak{S}_p(\mathcal{H})$ it follows from the Fredholm alternative (see, e.g. [78, Corollary to Theorem VI.14]) that $0 \in \rho(M(z))$ for all $z \in \mathbb{C}_+$. It is clear that

$$-M(z)^{-1} = D + L(z), \quad z \in \mathbb{C}_+, \quad (2.11)$$

holds with $L(z) := C^{-1} - M(z)^{-1}$, $z \in \mathbb{C}_+$ and the boundedly invertible self-adjoint operator $D := -C^{-1}$. Since

$$L(z) = C^{-1} - M(z)^{-1} = C^{-1}K(z)M(z)^{-1}, \quad z \in \mathbb{C}_+,$$

and $K(z) \in \mathfrak{S}_p(\mathcal{H})$, we conclude $L(z) \in \mathfrak{S}_p(\mathcal{H})$, $z \in \mathbb{C}_+$. Moreover, as C^{-1} is a bounded self-adjoint operator one gets

$$\operatorname{Im} L(z) = \operatorname{Im}(-M(z)^{-1}) = (M(z)^*)^{-1}(\operatorname{Im} K(z))M(z)^{-1}, \quad z \in \mathbb{C}_+,$$

where in the last equality we have used (2.10). As $K(\cdot) \in R^s[\mathcal{H}]$ by assumption we have $\ker(\operatorname{Im} K(z)) = \{0\}$ and this yields $\ker(\operatorname{Im} L(z)) = \{0\}$ for all $z \in \mathbb{C}_+$. We have shown that $L(\cdot) : \mathbb{C}_+ \rightarrow \mathfrak{S}_p(\mathcal{H})$ is a strict Nevanlinna function, $L(\cdot) \in R^s[\mathcal{H}]$, and hence it follows from (2.11) that $-M^{-1}(\cdot) \in R_{\mathfrak{S}_p}^{\text{reg}}[\mathcal{H}]$. \square

The assertions in the next lemma on the boundary values of \mathfrak{S}_1 -regular Nevanlinna functions follow from well-known results due to Birman and Èntina [25], de Branges [26], and Naboko [70]; cf. [44, Theorem 2.2].

Lemma 2.6. *Let $M(\cdot)$ be an \mathfrak{S}_1 -regular Nevanlinna function, $M(\cdot) \in R_{\mathfrak{S}_1}^{\text{reg}}[\mathcal{H}]$. Then the following assertions hold.*

- (i) $M(\lambda + i0) = \lim_{\varepsilon \rightarrow +0} M(\lambda + i\varepsilon)$ exists for a.e. $\lambda \in \mathbb{R}$ in the norm of $\mathcal{B}(\mathcal{H})$;
- (ii) $M(\lambda + i0)$ is boundedly invertible in \mathcal{H} for a.e. $\lambda \in \mathbb{R}$;
- (iii) $M(\lambda + i\varepsilon) - M(\lambda + i0) \in \mathfrak{S}_p(\mathcal{H})$ for $p \in (1, \infty]$, $\varepsilon > 0$ and a.e. $\lambda \in \mathbb{R}$, and

$$\lim_{\varepsilon \rightarrow +0} \|M(\lambda + i\varepsilon) - M(\lambda + i0)\|_{\mathfrak{S}_p(\mathcal{H})} = 0;$$

- (iv) $\operatorname{Im} M(\lambda + i0) = \lim_{\varepsilon \rightarrow +0} \operatorname{Im} M(\lambda + i\varepsilon)$ exists for a.e. $\lambda \in \mathbb{R}$ in the \mathfrak{S}_1 -norm.

Proof. By assumption there exists a Nevanlinna function $K(\cdot)$ with values in $\mathfrak{S}_1(\mathcal{H})$ such that $M(z) = C + K(z)$, $z \in \mathbb{C}_+$, holds with some bounded and boundedly invertible self-adjoint operator C . It follows from [25,26,70] (see, e.g. [44, Theorem 2.2]) that the limit $K(\lambda + i0)$ exists for a.e. $\lambda \in \mathbb{R}$ in the \mathfrak{S}_p -norm for all $p > 1$, and that the limit $\operatorname{Im} K(\lambda + i0)$ exists for a.e. $\lambda \in \mathbb{R}$ in the \mathfrak{S}_1 -norm. This yields assertions (i), (iii), and (iv).

In order to prove (ii) we recall that $-M(\cdot)^{-1}$ is \mathfrak{S}_1 -regular by Lemma 2.5 and hence the boundary values $M(\lambda + i0)^{-1}$ exist for a.e. $\lambda \in \mathbb{R}$ in the operator norm. Hence (ii) follows from the identity

$$M(\lambda + i\varepsilon)M(\lambda + i\varepsilon)^{-1} = M(\lambda + i\varepsilon)^{-1}M(\lambda + i\varepsilon) = I_{\mathcal{H}}, \quad \lambda \in \mathbb{R},$$

after passing to the limit $\varepsilon \rightarrow +0$ in the operator norm. \square

In the next lemma we investigate B -generalized boundary triples with \mathfrak{S}_p -regular Weyl functions. In particular, it turns out that the symmetric extension $A_1 = T \upharpoonright \ker(\Gamma_1)$ is self-adjoint and a Krein type resolvent formula is obtained; cf. [14,17,37,38].

Proposition 2.7. *Let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for S^* such that the corresponding Weyl function $M(\cdot)$ is \mathfrak{S}_p -regular for some $p \in (0, \infty]$. Then the following assertions hold.*

- (i) Π is a double B -generalized boundary triple for S^* ;
- (ii) The Weyl function corresponding to the transposed B -generalized boundary triple $\Pi^\top = \{\mathcal{H}, \Gamma_1, -\Gamma_0\}$ is \mathfrak{S}_p -regular;
- (iii) The operators A_0 and A_1 are \mathfrak{S}_p -resolvent comparable and

$$(A_1 - z)^{-1} - (A_0 - z)^{-1} = -\gamma(z)M(z)^{-1}\gamma(\bar{z})^* \in \mathfrak{S}_p(\mathfrak{H}) \quad (2.12)$$

holds for all $z \in \rho(A_0) \cap \rho(A_1)$.

Proof. (i) Since the Weyl function $M(\cdot)$ is \mathfrak{S}_p -regular by assumption, Lemma 2.5 implies, in particular, that $M(z)^{-1} \in \mathcal{B}(\mathcal{H})$ for all $z \in \mathbb{C} \setminus \mathbb{R}$. This yields

$$\operatorname{ran}(\Gamma_1) = \operatorname{ran}(M(z)) = \mathcal{H}.$$

Next we check that $A_1 = T \upharpoonright \ker(\Gamma_1)$ is self-adjoint in \mathfrak{H} . First of all it follows from the abstract Green's identity (2.1) that A_1 is symmetric. Let $z \in \mathbb{C} \setminus \mathbb{R}$, fix $f \in \mathfrak{H}$ and consider

$$h := (A_0 - z)^{-1}f - \gamma(z)M(z)^{-1}\gamma(\bar{z})^*f.$$

From Definition 2.3 and (2.6) we obtain

$$\Gamma_1 h = \Gamma_1(A_0 - z)^{-1}f - \Gamma_1\gamma(z)M(z)^{-1}\gamma(\bar{z})^*f = 0$$

and hence $h \in \text{dom}(A_1)$. Since $\text{ran } \gamma(z) \subset \ker(T - z)$ one gets

$$(A_1 - z)h = (T - z)((A_0 - z)^{-1}f - \gamma(z)M(z)^{-1}\gamma(\bar{z})^*f) = f$$

and we conclude the Krein type resolvent formula (2.12) in (iii) and $\text{ran}(A_1 - z) = \mathcal{H}$ for $z \in \mathbb{C} \setminus \mathbb{R}$. Hence the symmetric operator A_1 is self-adjoint in \mathfrak{H} and it follows that Π is a double B -generalized boundary triple for S^* .

(ii) The Weyl function corresponding to the transposed B -generalized boundary triple $\Pi^\top = \{\mathcal{H}, \Gamma_1, -\Gamma_0\}$ is given by

$$M^\top(z) = -M(z)^{-1}, \quad z \in \rho(A_0) \cap \rho(A_1), \quad (2.13)$$

which is \mathfrak{S}_p -regular by Lemma 2.5.

(iii) Since $M(\cdot)$ is \mathfrak{S}_p -regular it follows that $\text{Im } M(z) \in \mathfrak{S}_p(\mathcal{H})$ for $z \in \mathbb{C} \setminus \mathbb{R}$ and hence $\gamma(z)^*\gamma(z) \in \mathfrak{S}_p(\mathcal{H})$ by (2.8). This implies $\gamma(z) \in \mathfrak{S}_{2p}(\mathcal{H}, \mathfrak{H})$ and $\gamma(z)^* \in \mathfrak{S}_{2p}(\mathfrak{H}, \mathcal{H})$ for $z \in \mathbb{C} \setminus \mathbb{R}$, and the resolvent formula in (2.12) together with $0 \in \rho(M(z))$, $z \in \mathbb{C} \setminus \mathbb{R}$, yields the \mathfrak{S}_p -property of the resolvent difference in (2.12) for $z \in \mathbb{C} \setminus \mathbb{R}$, and hence for all $z \in \rho(A_0) \cap \rho(A_1)$. \square

Proposition 2.7 (iii) admits the following useful improvement.

Corollary 2.8. *Let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for S^* such that the corresponding Weyl function $M(\cdot)$ is \mathfrak{S}_∞ -regular and assume that $\text{Im } M(z) \in \mathfrak{S}_p(\mathcal{H})$ for some $p \in (0, \infty)$ and $z \in \mathbb{C}_+$. Then*

$$(A_1 - z)^{-1} - (A_0 - z)^{-1} \in \mathfrak{S}_p(\mathfrak{H}), \quad z \in \rho(A_0) \cap \rho(A_1). \quad (2.14)$$

Proof. The assumption $\text{Im } M(z) \in \mathfrak{S}_p(\mathcal{H})$ for some $p \in (0, \infty)$ and $z \in \mathbb{C}_+$ together with (2.8) yields $\gamma(z)^*\gamma(z) \in \mathfrak{S}_p(\mathcal{H})$, and hence $\gamma(z) \in \mathfrak{S}_{2p}(\mathcal{H}, \mathfrak{H})$. The Krein type formula in (2.12) implies (2.14) for $z \in \mathbb{C}_+$, and hence also for all $z \in \rho(A_0) \cap \rho(A_1)$. \square

Next we show that the p -resolvent comparability condition (2.12) guarantees the existence of a B -generalized boundary triple such that the corresponding Weyl function is \mathfrak{S}_p -regular.

Proposition 2.9. *Let A and B be self-adjoint operators in \mathfrak{H} and assume that the closed symmetric operator $S = A \cap B$ is densely defined. Then*

$$\text{dom}(A) + \text{dom}(B)$$

is dense in $\text{dom}(S^)$ with respect to the graph norm and the following assertions hold.*

(i) There is a B -generalized boundary triple $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ for S^* such that

$$A = T \upharpoonright \ker(\Gamma_0) = A_0 \quad \text{and} \quad B = T \upharpoonright \ker(\Gamma_1) = A_1. \quad (2.15)$$

(ii) If for some $z \in \mathbb{C} \setminus \mathbb{R}$ and some $p \in (0, \infty]$ the condition

$$(B - z)^{-1} - (A - z)^{-1} \in \mathfrak{S}_p(\mathfrak{H}) \quad (2.16)$$

is satisfied, then there exists a double B -generalized boundary triple $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ such that (2.15) holds and the corresponding Weyl function $M(\cdot)$ is \mathfrak{S}_p -regular.

Proof. In order to see that $\text{dom}(A) + \text{dom}(B)$ is dense in $\text{dom}(S^*)$ with respect to the graph norm assume that $h \in \text{dom}(S^*)$ is such that

$$(f_A + f_B, h) + (S^*(f_A + f_B), S^*h) = 0 \quad \text{for all } f_A \in \text{dom}(A), f_B \in \text{dom}(B).$$

Then $(Af_A, S^*h) = (f_A, -h)$ and $(Bf_B, S^*h) = (f_B, -h)$ for all $f_A \in \text{dom}(A)$ and $f_B \in \text{dom}(B)$ yield $S^*h \in \text{dom}(A) \cap \text{dom}(B) = \text{dom}(S)$ and $(I + SS^*)h = 0$. Since the operator $I + SS^*$ is uniformly positive one gets $h = 0$, that is, $\text{dom}(A) + \text{dom}(B)$ is dense in $\text{dom}(S^*)$ with respect to the graph norm.

(i) Observe first that $S = A \cap B$ is a densely defined, closed, symmetric operator with equal deficiency indices. Hence there exists an ordinary boundary triple $\Pi' = \{\mathcal{H}, \Gamma'_0, \Gamma'_1\}$ for S^* such that $B = S^* \upharpoonright \ker(\Gamma'_0)$; cf. [36, 38]. Furthermore, as A and B are disjoint self-adjoint extensions of S there exists a self-adjoint operator $\Theta = \Theta^* \in \mathcal{C}(\mathcal{H})$ such that

$$A = S^* \upharpoonright \text{dom}(A), \quad \text{dom}(A) = \ker(\Gamma'_1 - \Theta\Gamma'_0),$$

see e.g. [38, Proposition 1.4]. We consider the mappings

$$\Gamma_0 := \Gamma'_1 - \Theta\Gamma'_0 \quad \text{and} \quad \Gamma_1 := -\Gamma'_0$$

defined on

$$\text{dom}(\Gamma_0) = \text{dom}(\Gamma_1) := \text{dom}(A) + \text{dom}(B)$$

and set

$$T := S^* \upharpoonright \text{dom}(T), \quad \text{dom}(T) := \text{dom}(A) + \text{dom}(B).$$

We claim that $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is a B -generalized boundary triple for S^* such that (2.15) holds. Note first that $A = T \upharpoonright \ker(\Gamma_0) = A_0$, $B = T \upharpoonright \ker(\Gamma_1) = A_1$, and that A and B are disjoint self-adjoint extensions of S by construction. Therefore the argument in the beginning of the proof implies that $\text{dom}(T) = \text{dom}(A) + \text{dom}(B)$ is dense in $\text{dom}(S^*)$ equipped with the graph norm and hence $\overline{T} = S^*$. Moreover, since $\Theta = \Theta^*$ and the

abstract Green's identity (2.2) holds for the ordinary boundary triple Π' we obtain for $f, g \in \text{dom}(T)$

$$\begin{aligned} (\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g) &= (-\Gamma'_0 f, (\Gamma'_1 - \Theta \Gamma'_0)g) - ((\Gamma'_1 - \Theta \Gamma'_0)f, -\Gamma'_0 g) \\ &= (\Gamma'_1 f, \Gamma'_0 g) - (\Gamma'_0 f, \Gamma'_1 g) = (Tf, g) - (f, Tg), \end{aligned}$$

that is, the abstract Green's identity (2.1) holds. In order to verify $\text{ran}(\Gamma_0) = \mathcal{H}$ fix $h \in \mathcal{H}$. Since Π' is an ordinary boundary triple there exists $f_0 \in \text{dom}(B) = \ker(\Gamma'_0)$ such that $\Gamma'_1 f_0 = h$. We then obtain

$$\Gamma_0 f_0 = (\Gamma'_1 - \Theta \Gamma'_0) f_0 = \Gamma'_1 f_0 = h$$

and hence $\text{ran}(\Gamma_0) = \mathcal{H}$. Summing up, we have shown that Π is a B -generalized boundary triple such that (2.15) holds.

(ii) Now we choose an ordinary boundary triple $\Pi'' = \{\mathcal{H}, \Gamma''_0, \Gamma''_1\}$ for S^* such that $A = S^* \upharpoonright \ker(\Gamma''_0)$. Since A and B are disjoint extensions of S there exists an operator $\Theta = \Theta^* \in \mathcal{C}(\mathcal{H})$ such that

$$B = S^* \upharpoonright \text{dom}(B), \quad \text{dom}(B) = \ker(\Gamma''_1 - \Theta \Gamma''_0). \quad (2.17)$$

It follows from [37, Theorem 2] that the condition (2.16) is equivalent to the condition $(\Theta - \xi)^{-1} \in \mathfrak{S}_p(\mathcal{H})$ for all $\xi \in \rho(\Theta)$. In particular, $\rho(\Theta) \cap \mathbb{R} \neq \emptyset$, and in the following we assume without loss of generality that $0 \in \rho(\Theta)$. Denote the spectral function of the self-adjoint operator Θ by $E_\Theta(\cdot)$, let $\text{sgn}(\Theta) = \int_{\mathbb{R}} \text{sgn}(t) dE_\Theta(t)$ and recall the polar decomposition

$$\Theta = |\Theta|^{1/2} \text{sgn}(\Theta) |\Theta|^{1/2} = \text{sgn}(\Theta) |\Theta| = |\Theta| \text{sgn}(\Theta).$$

As $\Theta^{-1} \in \mathfrak{S}_p(\mathcal{H})$ we have $|\Theta|^{-1/2} \in \mathfrak{S}_{2p}(\mathcal{H})$ and $\ker(|\Theta|^{-1/2}) = \{0\}$. We consider the mappings

$$\Gamma_0 := |\Theta|^{1/2} \Gamma''_0 \quad \text{and} \quad \Gamma_1 := |\Theta|^{-1/2} (\Gamma''_1 - \Theta \Gamma''_0) \quad (2.18)$$

defined on

$$\text{dom}(\Gamma_0) = \text{dom}(\Gamma_1) := \{f \in \text{dom}(S^*) : \Gamma''_0 f \in \text{dom}(|\Theta|^{1/2})\}. \quad (2.19)$$

We set

$$T := S^* \upharpoonright \text{dom}(T), \quad \text{dom}(T) := \text{dom}(\Gamma_0) = \text{dom}(\Gamma_1),$$

and we claim that $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is a double B -generalized boundary triple for S^* . First of all we have for $f, g \in \text{dom}(T)$

$$\begin{aligned}
& (\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g) \\
&= (|\Theta|^{-1/2}(\Gamma_1'' - \Theta \Gamma_0'')f, |\Theta|^{1/2} \Gamma_0'' g) - (|\Theta|^{1/2} \Gamma_0'' f, |\Theta|^{-1/2}(\Gamma_1'' - \Theta \Gamma_0'')g) \\
&= ((\Gamma_1'' - \Theta \Gamma_0'')f, \Gamma_0'' g) - (\Gamma_0'' f, (\Gamma_1'' - \Theta \Gamma_0'')g) \\
&= (\Gamma_1'' f, \Gamma_0'' g) - (\Gamma_0'' f, \Gamma_1'' g)
\end{aligned}$$

and since Π'' is an ordinary boundary triple the abstract Green's identity (2.1) follows. The condition $\text{ran}(\Gamma_0) = \mathcal{H}$ is satisfied since $0 \in \rho(\Theta)$, and thus also $0 \in \rho(|\Theta|^{1/2})$. It is also clear from the definition of Γ_0 in (2.18)–(2.19) that

$$\ker(\Gamma_0) = \ker(\Gamma_0'') = \text{dom}(A). \quad (2.20)$$

Next it will be shown that

$$\ker(\Gamma_1) = \text{dom}(B) \quad (2.21)$$

holds. In fact, the inclusion $\ker(\Gamma_1) \subset \text{dom}(B)$ in (2.21) follows from the definition of Γ_1 in (2.18)–(2.19) and $\ker(|\Theta|^{-1/2}) = \{0\}$. For the remaining inclusion let $f \in \text{dom}(B)$. Then $\Gamma_1'' f = \Theta \Gamma_0'' f$ by (2.17) and, in particular,

$$\Gamma_0'' f \in \text{dom}(\Theta) \subset \text{dom}(|\Theta|^{1/2}).$$

Hence $\text{dom}(B) \subset \text{dom}(T)$ and $\Gamma_1 f = 0$ is clear, that is, $\text{dom}(B) \subset \ker(\Gamma_1)$ and thus (2.21) is shown. Combining (2.20) with (2.21) yields (2.15). Moreover, we have $\overline{T} = S^*$ since

$$\text{dom}(A) + \text{dom}(B) = \ker(\Gamma_0) + \ker(\Gamma_1) \subset \text{dom}(T)$$

and $\text{dom}(A) + \text{dom}(B)$ is dense in $\text{dom}(S^*)$ equipped with the graph norm (as A and B are disjoint self-adjoint extensions of S). Summing up, we have shown that $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is a B -generalized boundary triple for S^* such that (2.15) holds.

It remains to verify that the Weyl function corresponding to Π is \mathfrak{S}_p -regular; Proposition 2.7 (i) then implies that Π is a double B -generalized boundary triple. For this denote the Weyl function corresponding to the ordinary boundary triple Π'' by $M''(\cdot)$ and recall that $M''(z)\Gamma_0'' f_z = \Gamma_1'' f_z$ for $f_z \in \ker(S^* - z)$ and $z \in \rho(A)$. We claim that the Weyl function corresponding to Π is given by

$$M(z) = |\Theta|^{-1/2} M''(z) |\Theta|^{-1/2} - \text{sgn}(\Theta), \quad z \in \rho(A). \quad (2.22)$$

In fact, for $f_z \in \ker(T - z)$ we compute

$$\begin{aligned}
& (|\Theta|^{-1/2}M''(z)|\Theta|^{-1/2} - \operatorname{sgn}(\Theta))\Gamma_0 f_z \\
&= |\Theta|^{-1/2}M''(z)\Gamma_0'' f_z - \operatorname{sgn}(\Theta)|\Theta|^{1/2}\Gamma_0'' f_z \\
&= |\Theta|^{-1/2}(\Gamma_1'' f_z - |\Theta|^{1/2}\operatorname{sgn}(\Theta)|\Theta|^{1/2}\Gamma_0'' f_z) \\
&= |\Theta|^{-1/2}(\Gamma_1'' f_z - \Theta\Gamma_0'' f_z) = \Gamma_1 f_z
\end{aligned}$$

and hence (2.22) follows by Definition 2.3. Let $K(z) := |\Theta|^{-1/2}M''(z)|\Theta|^{-1/2}$, $z \in \mathbb{C}_+$ and let $C := -\operatorname{sgn}(\Theta)$. Note that C is a boundedly invertible self-adjoint operator and that $|\Theta|^{-1/2} \in \mathfrak{S}_{2p}(\mathcal{H})$ and $M''(z) \in \mathcal{B}(\mathcal{H})$ yield $K(z) \in \mathfrak{S}_p(\mathcal{H})$, $z \in \mathbb{C}_+$. Moreover, as $M''(\cdot) \in R^u[\mathcal{H}]$ it follows that $K(\cdot) \in R^s[\mathcal{H}]$, and hence the Weyl function $M(\cdot)$ is \mathfrak{S}_p -regular. \square

In applications to scattering problems it is important to know whether the resolvent p -comparability condition (2.12), (2.16) yields the \mathfrak{S}_p -regularity of the Weyl function. A converse statement to Proposition 2.7 is false for arbitrary double B -generalized boundary triples, while Proposition 2.9 ensures the existence of such a double B -generalized boundary triple. However in the following proposition we present an affirmative answer to this question under certain additional explicit assumptions.

Proposition 2.10. *Let A and B be self-adjoint operators in \mathfrak{H} such that*

$$R_{B,A}(z) := (B - z)^{-1} - (A - z)^{-1} \in \mathfrak{S}_p(\mathfrak{H}) \quad (2.23)$$

for some $z \in \mathbb{C} \setminus \mathbb{R}$ and some $p \in (0, \infty]$, and assume that the closed symmetric operator $S = A \cap B$ is densely defined. Assume, in addition, that there exists $\lambda_0 \in \rho(A) \cap \rho(B) \cap \mathbb{R}$ such that

$$\pm R_{B,A}(\lambda_0) \geq 0. \quad (2.24)$$

If $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ is a double B -generalized boundary triple for S^* such that condition (2.15) holds then the corresponding Weyl function $M(\cdot)$ is \mathfrak{S}_p -regular.

Proof. Since Π is a double B -generalized boundary triple, the values of the Weyl function $M(\cdot)$ and the function $-M(\cdot)^{-1}$ are in $\mathcal{B}(\mathcal{H})$. Moreover, the assumption $\lambda_0 \in \rho(A) \cap \rho(B) \cap \mathbb{R}$ ensures that $-M(\lambda_0)^{-1} \in \mathcal{B}(\mathcal{H})$ is a self-adjoint operator and we have

$$R_{B,A}(\lambda_0) = (B - \lambda_0)^{-1} - (A - \lambda_0)^{-1} = -\gamma(\lambda_0)M(\lambda_0)^{-1}\gamma(\lambda_0)^* \quad (2.25)$$

by Proposition 2.7 (iii). Assume that $R_{A,B}(\lambda_0) \geq 0$ in (2.24). Then by (2.25)

$$(R_{A,B}(\lambda_0)f, f) = (-M(\lambda_0)^{-1}\gamma(\lambda_0)^*f, \gamma(\lambda_0)^*f) \geq 0, \quad f \in \mathfrak{H},$$

and since $\operatorname{ran}(\gamma(\lambda_0)^*)$ is dense in \mathcal{H} (see Section 2.1) we have $-M(\lambda_0)^{-1} \geq 0$. Setting $T(\lambda_0) := \gamma(\lambda_0)(-M(\lambda_0))^{-1/2} \in \mathcal{B}(\mathcal{H}, \mathfrak{H})$ and using the assumption (2.23) for some, and hence for all, $z \in \rho(A) \cap \rho(B)$ we conclude from (2.25) that

$$R_{B,A}(\lambda_0) = T(\lambda_0)T(\lambda_0)^* \in \mathfrak{S}_p(\mathfrak{H}).$$

This relation yields $T(\lambda_0)^* \in \mathfrak{S}_{2p}(\mathfrak{H}, \mathcal{H})$ and $T(\lambda_0) \in \mathfrak{S}_{2p}(\mathcal{H}, \mathfrak{H})$, and hence $\gamma(\lambda_0) = T(\lambda_0)(-M(\lambda_0))^{1/2} \in \mathfrak{S}_{2p}(\mathcal{H}, \mathfrak{H})$. It then follows from (2.5) that

$$\gamma(z) \in \mathfrak{S}_{2p}(\mathcal{H}, \mathfrak{H}) \quad \text{and} \quad \gamma(\xi)^* \in \mathfrak{S}_{2p}(\mathfrak{H}, \mathcal{H}), \quad z, \xi \in \rho(A).$$

Combining this with (2.7) it follows that $M(z) - M(\lambda_0) \in \mathfrak{S}_p(\mathcal{H})$. Therefore, setting $C := M(\lambda_0)$ and $K(z) := M(z) - M(\lambda_0)$, $z \in \mathbb{C}_+$, we arrive at the representation (2.9). Note that $C = M(\lambda_0)$ is a boundedly invertible self-adjoint operator. Furthermore, since $\operatorname{Im} K(z) = \operatorname{Im} M(z)$ and $M(\cdot) \in R^s[\mathcal{H}]$ we conclude $K(\cdot) \in R^s[\mathcal{H}]$, that is, the Weyl function $M(\cdot)$ is \mathfrak{S}_p -regular. \square

Remark 2.11. Condition (2.24) is satisfied if the symmetric operator $S = A \cap B$ is semibounded from below and A is chosen to be its Friedrichs extension. In this case (2.23) yields the semiboundedness of the operator B and the inequality (2.24) holds for any λ_0 smaller than the lower bound of B .

Remark 2.12. The density of $\operatorname{dom}(A) + \operatorname{dom}(B)$ in \mathfrak{H} under the conditions of Proposition 2.9 is well known (see for instance [36]). The simple proof presented here and which does not exploit the second Neumann formula seems to be new.

Remark 2.13. Proposition 2.7 (i) can also be viewed as an immediate consequence from the fact that the values of $M^{-1}(\cdot)$ are in $\mathcal{B}(\mathcal{H})$; cf. [34,38]. For the convenience of the reader we have presented a simple direct proof.

In the proofs of the results in Sections 4 and 5 we shall occasionally make use of the following lemma.

Lemma 2.14. Let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for $\overline{T} = S^*$ and let $M(\cdot)$ be the corresponding Weyl function. Assume that $A_1 = A_1^*$ and that $\xi \in \rho(A_0)$. Then the following equivalence holds:

$$\xi \in \sigma_p(A_1) \Leftrightarrow 0 \in \sigma_p(M(\xi)).$$

Proof. Assume first that $0 \in \sigma_p(M(\xi))$. Then there exists $\psi \in \mathcal{H}$, $\psi \neq 0$, such that $M(\xi)\psi = 0$. Since $\operatorname{ran}(\Gamma_0) = \mathcal{H}$ one finds $f_\xi \in \ker(T - \xi)$, $f_\xi \neq 0$, with $\psi = \Gamma_0 f_\xi$. Then $\Gamma_1 f_\xi = M(\xi)\psi = 0$ and $f_\xi \in \operatorname{dom}(A_1)$. This shows $f_\xi \in \ker(A_1 - \xi)$ and $\xi \in \sigma_p(A_1)$.

Conversely, assume that $f_\xi \in \ker(A_1 - \xi)$, $f_\xi \neq 0$. Then $\Gamma_1 f_\xi = 0$ and $\psi := \Gamma_0 f_\xi \neq 0$ since otherwise $\xi \in \sigma_p(A_0)$. Then $M(\xi)\psi = M(\xi)\Gamma_0 f_\xi = \Gamma_1 f_\xi = 0$ and hence it follows that $0 \in \sigma_p(M(\xi))$. \square

3. A representation of the scattering matrix

Let A and B be self-adjoint operators in a Hilbert space \mathfrak{H} and assume that they are resolvent comparable, i.e. their resolvent difference is a trace class operator,

$$(B - i)^{-1} - (A - i)^{-1} \in \mathfrak{S}_1(\mathfrak{H}). \quad (3.1)$$

Denote by $\mathfrak{H}^{ac}(A)$ the absolutely continuous subspace of A and let $P^{ac}(A)$ be the orthogonal projection in \mathfrak{H} onto $\mathfrak{H}^{ac}(A)$. In accordance with the Birman–Krein theorem, under the assumption (3.1) the *wave operators*

$$W_+(A, B) := s - \lim_{t \rightarrow +\infty} e^{itB} e^{-itA} P^{ac}(A)$$

and

$$W_-(A, B) := s - \lim_{t \rightarrow -\infty} e^{itB} e^{-itA} P^{ac}(A)$$

exist and are complete, i.e. the ranges of $W_+(A, B)$ and $W_-(A, B)$ coincide with the absolutely continuous subspace $\mathfrak{H}^{ac}(B)$ of B ; cf. [12,59,79,81,82]. The *scattering operator* $S(A, B)$ of the *scattering system* is defined by

$$S(A, B) = W_+(A, B)^* W_-(A, B).$$

The operator $S(A, B)$ commutes with A and is unitary in $\mathfrak{H}^{ac}(A)$, hence it is unitarily equivalent to a multiplication operator induced by a family $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ of unitary operators in a spectral representation of the absolutely continuous part A^{ac} of A ,

$$A^{ac} := A \upharpoonright \text{dom}(A) \cap \mathfrak{H}^{ac}(A).$$

The family $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ is called the *scattering matrix* of the scattering system $\{A, B\}$.

In Theorem 3.1 and Corollary 3.3 below we shall provide a representation of the scattering matrix $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ of the system $\{A, B\}$ in an extension theory framework using B -generalized boundary triples and their Weyl functions. It is assumed that the closed symmetric operator $S = A \cap B$ is densely defined; in the more general framework of non-densely defined symmetric operators this assumption can be dropped. First we discuss the case that $S = A \cap B$ is simple, i.e. S does not contain a self-adjoint part or, equivalently, the condition

$$\mathfrak{H} = \text{clsp}\{\ker(S^* - z) : z \in \mathbb{C} \setminus \mathbb{R}\}$$

is satisfied; cf. [60]. In the sequel the abbreviation a.e. means “almost everywhere with respect to the Lebesgue measure”.

Theorem 3.1. *Let A and B be self-adjoint operators in a Hilbert space \mathfrak{H} , assume that the closed symmetric operator $S = A \cap B$ is densely defined and simple, and let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for S^* such that $A = T \upharpoonright \ker(\Gamma_0)$ and $B = T \upharpoonright \ker(\Gamma_1)$. Assume, in addition, that the Weyl function $M(\cdot)$ corresponding to Π is \mathfrak{S}_1 -regular.*

Then $\{A, B\}$ is a complete scattering system and

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda), \quad \mathcal{H}_\lambda := \overline{\text{ran}(\text{Im } M(\lambda + i0))},$$

forms a spectral representation of A^{ac} such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A, B\}$ admits the representation

$$S(A, B; \lambda) = I_{\mathcal{H}_\lambda} - 2i\sqrt{\text{Im } M(\lambda + i0)} M(\lambda + i0)^{-1} \sqrt{\text{Im } M(\lambda + i0)}.$$

Proof. The proof of Theorem 3.1 consists of three separate steps and is essentially based on Theorem A.2. Parts of the proof follow the lines in [20, Proof of Theorem 3.1], where the special case of a symmetric operator S with finite deficiency indices was treated.

First of all we note that the \mathfrak{S}_1 -regularity assumption on $M(\cdot)$ together with Proposition 2.7 (iii) ensures that the resolvent difference of A and B is a trace class operator. Hence the wave operators $W_\pm(A, B)$ exist and are complete and $\{A, B\}$ is a complete scattering system, see, e.g. [82, Theorem VI.5.1].

Step 1. According to Proposition 2.7 (iii) the resolvent difference of A and B in (3.1) can be written in a Krein type resolvent formula of the form

$$(B - z)^{-1} - (A - z)^{-1} = -\gamma(z)M(z)^{-1}\gamma(\bar{z})^*, \quad z \in \rho(A) \cap \rho(B). \quad (3.2)$$

In particular, from (3.2) and (2.5) we get

$$\begin{aligned} (B - i)^{-1} - (A - i)^{-1} &= -\gamma(i)M(i)^{-1}\gamma(-i)^* \\ &= -(A + i)(A - i)^{-1}\gamma(-i)M(i)^{-1}\gamma(-i)^* = \phi(A)CGC^* \end{aligned}$$

where

$$\phi(t) := \frac{t+i}{t-i}, \quad t \in \mathbb{R}, \quad C := \gamma(-i) \quad \text{and} \quad G := -M(i)^{-1}. \quad (3.3)$$

We claim that the condition

$$\mathfrak{H}^{ac}(A) = \text{clsp} \{E_A^{ac}(\delta) \text{ran } C : \delta \in \mathfrak{B}(\mathbb{R})\} \quad (3.4)$$

in Theorem A.2 is satisfied. In fact, since S is assumed to be simple we have

$$\mathfrak{H} = \text{clsp} \{\ker(S^* - z) : z \in \mathbb{C} \setminus \mathbb{R}\}.$$

Furthermore, using $\ker(S^* - z) = \overline{\ker(T - z)}$, $z \in \mathbb{C} \setminus \mathbb{R}$, which follows from (2.4), and $\operatorname{ran}(\gamma(z)) = \ker(T - z)$, $z \in \mathbb{C} \setminus \mathbb{R}$, it follows that

$$\begin{aligned}\mathfrak{H} &= \operatorname{clsp} \{ \ker(T - z) : z \in \mathbb{C} \setminus \mathbb{R} \} \\ &= \operatorname{clsp} \{ \gamma(z)h : z \in \mathbb{C} \setminus \mathbb{R}, h \in \mathcal{H} \} \\ &= \operatorname{clsp} \{ (A + i)(A - z)^{-1}\gamma(-i)h : z \in \mathbb{C} \setminus \mathbb{R}, h \in \mathcal{H} \} \\ &= \operatorname{clsp} \{ (A + i)(A - z)^{-1}Ch : z \in \mathbb{C} \setminus \mathbb{R}, h \in \mathcal{H} \} \\ &= \operatorname{clsp} \{ E_A(\delta)Ch : h \in \mathcal{H}, \delta \in \mathfrak{B}(\mathbb{R}) \}\end{aligned}$$

and hence

$$\mathfrak{H}^{ac}(A) = \operatorname{clsp} \{ P^{ac}(A)E_A(\delta)Ch : h \in \mathcal{H}, \delta \in \mathfrak{B}(\mathbb{R}) \}.$$

Since $E_A^{ac}(\delta) = P^{ac}(A)E_A(\delta)$ this implies (3.4).

Step 2. Now we apply Theorem A.2 to obtain a preliminary form of the scattering matrix $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$. Since $M(\cdot)$ is \mathfrak{S}_1 -regular by assumption we have

$$\operatorname{Im} M(i) = \gamma(i)^* \gamma(i) \in \mathfrak{S}_1(\mathcal{H})$$

(see (2.8)) and hence $\gamma(i) \in \mathfrak{S}_2(\mathcal{H}, \mathfrak{H})$ and

$$C = \gamma(-i) = (I - 2i(A + i)^{-1})\gamma(i) \in \mathfrak{S}_2(\mathcal{H}, \mathfrak{H}).$$

Therefore the function $\lambda \mapsto C^* E_A((-\infty, \lambda))C$ is $\mathfrak{S}_1(\mathcal{H})$ -valued and in accordance with [25, Lemma 2.2] this function is $\mathfrak{S}_1(\mathcal{H})$ -differentiable for a.e. $\lambda \in \mathbb{R}$. We compute its derivative

$$\lambda \mapsto K(\lambda) = \frac{d}{d\lambda} C^* E_A((-\infty, \lambda))C$$

and the square root $\lambda \mapsto \sqrt{K(\lambda)}$ for a.e. $\lambda \in \mathbb{R}$. First we note that by the $\mathfrak{S}_1(\mathcal{H})$ -generalization of the Fatou theorem (see [25, Lemma 2.4])

$$\begin{aligned}K(\lambda) &= \lim_{\varepsilon \rightarrow 0+} \frac{1}{2\pi i} C^* ((A - \lambda - i\varepsilon)^{-1} - (A - \lambda + i\varepsilon)^{-1})C \\ &= \lim_{\varepsilon \rightarrow 0+} \frac{\varepsilon}{\pi} C^* ((A - \lambda - i\varepsilon)^{-1}(A - \lambda + i\varepsilon)^{-1})C\end{aligned}\tag{3.5}$$

for a.e. $\lambda \in \mathbb{R}$. On the other hand, inserting the formula

$$\gamma(\lambda + i\varepsilon) = (A + i)(A - \lambda - i\varepsilon)^{-1}\gamma(-i) = (A + i)(A - \lambda - i\varepsilon)^{-1}C$$

(see (2.5)) into (2.8) leads to

$$\begin{aligned}\operatorname{Im} M(\lambda + i\varepsilon) &= \varepsilon \gamma(\lambda + i\varepsilon)^* \gamma(\lambda + i\varepsilon) \\ &= \varepsilon C^*(I + A^2)(A - \lambda + i\varepsilon)^{-1}(A - \lambda - i\varepsilon)^{-1}C.\end{aligned}$$

Combining this relation with (3.5) we conclude

$$\operatorname{Im} M(\lambda + i0) = \lim_{\varepsilon \rightarrow 0+} \operatorname{Im} M(\lambda + i\varepsilon) = \pi(1 + \lambda^2)K(\lambda)$$

for a.e. $\lambda \in \mathbb{R}$. In particular, $\operatorname{ran}(\operatorname{Im} M(\lambda + i0)) = \operatorname{ran}(K(\lambda))$ for a.e. $\lambda \in \mathbb{R}$ and hence

$$\mathcal{H}_\lambda = \overline{\operatorname{ran}(\operatorname{Im} M(\lambda + i0))} = \overline{\operatorname{ran}(K(\lambda))} \quad \text{for a.e. } \lambda \in \mathbb{R}.$$

Therefore $L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$ is a spectral representation of A^{ac} and in accordance with Theorem A.2 the scattering matrix $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ is given by

$$\begin{aligned}S(A, B; \lambda) &= I_{\mathcal{H}_\lambda} + 2\pi i(1 + \lambda^2)^2 \sqrt{K(\lambda)} Z(\lambda) \sqrt{K(\lambda)} \\ &= I_{\mathcal{H}_\lambda} + 2i(1 + \lambda^2) \sqrt{\operatorname{Im} M(\lambda + i0)} Z(\lambda) \sqrt{\operatorname{Im} M(\lambda + i0)}\end{aligned}\tag{3.6}$$

for a.e. $\lambda \in \mathbb{R}$, where $Z(\cdot)$ is given by (A.6),

$$Z(\lambda) = \frac{1}{\lambda + i} Q^* Q + \frac{1}{(\lambda + i)^2} \phi(\lambda) G + \lim_{\varepsilon \rightarrow 0+} Q^* (B - (\lambda + i\varepsilon))^{-1} Q,\tag{3.7}$$

and

$$Q = \phi(A)CG = -(A + i)(A - i)^{-1} \gamma(-i) M(i)^{-1} = -\gamma(i) M(i)^{-1} \in \mathfrak{S}_2(\mathcal{H}, \mathfrak{H}).$$

Observe that due to the last inclusion the limit in (3.7) exists for a.e. $\lambda \in \mathbb{R}$ in every \mathfrak{S}_p -norm with $p > 1$ and the operator-valued function $Z(\cdot)$ in (3.7) is well defined a.e. on \mathbb{R} ; cf. Lemma 2.6.

Step 3. In the third and final step we prove that

$$Z(\lambda) = -\frac{1}{1 + \lambda^2} M(\lambda + i0)^{-1}\tag{3.8}$$

for a.e. $\lambda \in \mathbb{R}$. Then inserting this expression in (3.6) one arrives at the asserted form of the scattering matrix.

Applying the mapping Γ_0 to (3.2) and using $\ker(\Gamma_0) = \operatorname{dom}(A)$ and Definition 2.3 one gets

$$\Gamma_0(B - z)^{-1} = \Gamma_0(A - z)^{-1} - \Gamma_0 \gamma(z) M(z)^{-1} \gamma(\bar{z})^* = -M(z)^{-1} \gamma(\bar{z})^*\tag{3.9}$$

for $z \in \rho(A) \cap \rho(B)$ and hence

$$\Gamma_0(B + i)^{-1} = -M(-i)^{-1} \gamma(i)^* = (-\gamma(i) M(i)^{-1})^* = Q^*.$$

This yields

$$\begin{aligned} Q^*(B-z)^{-1}Q &= \Gamma_0(B+i)^{-1}(B-z)^{-1}Q \\ &= \Gamma_0(Q^*(B-\bar{z})^{-1}(B-i)^{-1})^* \\ &= \Gamma_0(\Gamma_0(B+i)^{-1}(B-\bar{z})^{-1}(B-i)^{-1})^*. \end{aligned} \quad (3.10)$$

In order to compute this expression we note that

$$\begin{aligned} (B+i)^{-1}(B-\bar{z})^{-1}(B-i)^{-1} \\ = \frac{-1}{1+\bar{z}^2}((B+i)^{-1} - (B-\bar{z})^{-1}) + \frac{1}{2i(\bar{z}-i)}((B+i)^{-1} - (B-i)^{-1}) \end{aligned}$$

and hence (3.9) implies

$$\begin{aligned} \Gamma_0(B+i)^{-1}(B-\bar{z})^{-1}(B-i)^{-1} &= \frac{1}{1+\bar{z}^2}(M(-i)^{-1}\gamma(i)^* - M(\bar{z})^{-1}\gamma(z)^*) \\ &\quad - \frac{1}{2i(\bar{z}-i)}(M(-i)^{-1}\gamma(i)^* - M(i)^{-1}\gamma(-i)^*). \end{aligned}$$

Taking into account that $(M(\bar{\mu})^{-1})^* = M(\mu)^{-1}$ for $\mu \in \rho(A) \cap \rho(B)$ we obtain for the adjoint

$$\begin{aligned} (\Gamma_0(B+i)^{-1}(B-\bar{z})^{-1}(B-i)^{-1})^* &= \frac{1}{1+z^2}(\gamma(i)M(i)^{-1} - \gamma(z)M(z)^{-1}) \\ &\quad + \frac{1}{2i(z+i)}(\gamma(i)M(i)^{-1} - \gamma(-i)M(-i)^{-1}). \end{aligned}$$

In turn, combining this identity with (3.10) yields

$$\begin{aligned} Q^*(B-z)^{-1}Qh &= \Gamma_0(\Gamma_0(B+i)^{-1}(B-\bar{z})^{-1}(B-i)^{-1})^* \\ &= \frac{1}{1+z^2}(M(i)^{-1} - M(z)^{-1}) + \frac{1}{2i(z+i)}(M(i)^{-1} - M(-i)^{-1}) \end{aligned}$$

for $z \in \rho(A) \cap \rho(B)$. Setting here $z = \lambda + i\varepsilon \in \mathbb{C}_+$ and passing to the limit as $\varepsilon \rightarrow 0$ one derives

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} Q^*(B - (\lambda + i\varepsilon))^{-1}Q &= \frac{1}{1+\lambda^2}(M(i)^{-1} - M(\lambda + i0)^{-1}) \\ &\quad + \frac{1}{2i(\lambda + i)}(M(i)^{-1} - M(-i)^{-1}) \end{aligned} \quad (3.11)$$

for a.e. $\lambda \in \mathbb{R}$; note that by Lemma 2.6 the limit $M(\lambda + i0)^{-1} \in \mathcal{B}(\mathcal{H})$ exists for a.e. $\lambda \in \mathbb{R}$.

Moreover, we have

$$\begin{aligned} Q^*Q &= (\gamma(i)M(i)^{-1})^*\gamma(i)M(i)^{-1} = M(-i)^{-1}\gamma(i)^*\gamma(i)M(i)^{-1} \\ &= \frac{1}{2i}M(-i)^{-1}(M(i) - M(-i))M(i)^{-1} = \frac{1}{2i}(M(-i)^{-1} - M(i)^{-1}). \end{aligned}$$

Inserting this relation and (3.11) into (3.7) and taking notations (3.3) into account we obtain for a.e. $\lambda \in \mathbb{R}$

$$\begin{aligned} Z(\lambda) &= \frac{1}{\lambda+i}Q^*Q + \frac{1}{(\lambda+i)^2}\phi(\lambda)G + Q^*(B - (\lambda+i0))^{-1}Q \\ &= \frac{1}{2i(\lambda+i)}(M(-i)^{-1} - M(i)^{-1}) - \frac{1}{1+\lambda^2}M(i)^{-1} \\ &\quad + \frac{1}{1+\lambda^2}(M(i)^{-1} - M(\lambda+i0)^{-1}) + \frac{1}{2i(\lambda+i)}(M(i)^{-1} - M(-i)^{-1}) \\ &= -\frac{1}{1+\lambda^2}M(\lambda+i0)^{-1}, \end{aligned}$$

that is, (3.8) holds. \square

Remark 3.2. Instead of the assumption that the Weyl function is \mathfrak{S}_1 -regular one may assume in Theorem 3.1 that $R_{B,A}(z) = (B-z)^{-1} - (A-z)^{-1} \in \mathfrak{S}_1(\mathfrak{H})$ holds for some $z \in \rho(A) \cap \rho(B)$ and $R_{B,A}(\lambda_0) \geq 0$ for some $\lambda_0 \in \mathbb{R} \cap \rho(A) \cap \rho(B)$; cf. Proposition 2.10.

Our next task is to drop the assumption of the simplicity of S in Theorem 3.1. If $S = A \cap B$ is not simple then the Hilbert space \mathfrak{H} admits an orthogonal decomposition $\mathfrak{H} = \mathfrak{H}_0 \oplus \mathfrak{H}'$ with $\mathfrak{H}_0 \neq \{0\}$ such that

$$S = S_0 \oplus S', \quad (3.12)$$

where S_0 is a self-adjoint operator in the Hilbert space \mathfrak{H}_0 and S' is a simple symmetric operator in the Hilbert space \mathfrak{H}' ; cf. [60]. It follows that there exist self-adjoint extensions A' and B' of S' in \mathfrak{H}' such that

$$A = S_0 \oplus A' \quad \text{and} \quad B = S_0 \oplus B'.$$

By restricting the boundary maps of a B -generalized boundary triple for S^* one obtains a B -generalized boundary triple for the operator $(S')^*$ with the same Weyl function. Applying Theorem 3.1 to the pair $\{A', B'\}$ yields the following variant of Theorem 3.1; cf. [20, Proof of Theorem 3.2] for the same argument in the special case of finite rank perturbations.

Corollary 3.3. *Let A and B be self-adjoint operators in a Hilbert space \mathfrak{H} , assume that the closed symmetric operator $S = A \cap B$ is densely defined and decomposed in $S = S_0 \oplus S'$ as*

in (3.12), and let $L^2(\mathbb{R}, d\lambda, \mathcal{G}_\lambda)$ be a spectral representation of S_0^{ac} . Let $\Pi = \{\mathcal{H}, \Gamma_0, \Gamma_1\}$ be a B -generalized boundary triple for S^* as in Theorem 3.1 such that the corresponding Weyl function $M(\cdot)$ is \mathfrak{S}_1 -regular.

Then $\{A, B\}$ is a complete scattering system and

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda \oplus \mathcal{G}_\lambda), \quad \mathcal{H}_\lambda := \overline{\text{ran}(\text{Im } M(\lambda + i0))},$$

forms a spectral representation of A^{ac} such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A, B\}$ admits the representation

$$S(A, B; \lambda) = \begin{pmatrix} S(A', B'; \lambda) & 0 \\ 0 & I_{\mathcal{G}_\lambda} \end{pmatrix},$$

where

$$S(A', B'; \lambda) = I_{\mathcal{H}_\lambda} - 2i\sqrt{\text{Im } M(\lambda + i0)} M(\lambda + i0)^{-1} \sqrt{\text{Im } M(\lambda + i0)}.$$

4. Scattering matrices for Schrödinger operators on exterior domains

Our main objective in this section is to derive representations of the scattering matrices for pairs of self-adjoint Schrödinger operators with Dirichlet, Neumann and Robin boundary conditions on unbounded domains with smooth compact boundaries in terms of Dirichlet-to-Neumann and Neumann-to-Dirichlet maps. After some necessary preliminaries in Sections 4.1 and 4.2 we formulate and prove our main results Theorem 4.3 and Theorem 4.7 in Sections 4.3 and 4.4, respectively. Both theorems follow in a similar way from our general result Theorem 3.1 by fixing a suitable B -generalized boundary triple and verifying that the corresponding Weyl function is \mathfrak{S}_1 -regular. We also mention that along the way we obtain classical results on singular value estimates of resolvent differences due to Birman, Grubb and others without any extra efforts; cf. Remarks 4.4 and 4.8.

4.1. Preliminaries on Sobolev spaces, trace maps, and Green's second identity

Let $\Omega \subset \mathbb{R}^n$ be an exterior domain, that is, $\mathbb{R}^n \setminus \Omega$ is bounded and closed, and assume that the boundary $\partial\Omega$ of Ω is C^∞ -smooth. We denote by $H^s(\Omega)$, $s \in \mathbb{R}$, the usual L^2 -based Sobolev spaces on the unbounded exterior domain Ω , and by $H^r(\partial\Omega)$, $r \in \mathbb{R}$, the corresponding Sobolev spaces on the compact C^∞ -boundary $\partial\Omega$. The corresponding scalar products will be denoted by (\cdot, \cdot) , and sometimes the space is used as an index.

Recall that the Dirichlet and Neumann trace operators γ_D and γ_N , originally defined as linear mappings from $C_0^\infty(\bar{\Omega})$ to $C^\infty(\partial\Omega)$, admit continuous extensions onto $H^2(\Omega)$ such that the mapping

$$\begin{pmatrix} \gamma_D \\ \gamma_N \end{pmatrix} : H^2(\Omega) \rightarrow H^{3/2}(\partial\Omega) \times H^{1/2}(\partial\Omega) \quad (4.1)$$

is surjective. The spaces

$$H_{\Delta}^s(\Omega) = \{f \in H^s(\Omega) : \Delta f \in L^2(\Omega)\}, \quad s \in [0, 2], \quad (4.2)$$

equipped with the Hilbert scalar products

$$(f, g)_{H_{\Delta}^s(\Omega)} = (f, g)_{H^s(\Omega)} + (\Delta f, \Delta g)_{L^2(\Omega)}, \quad f, g \in H_{\Delta}^s(\Omega), \quad (4.3)$$

will play an important role. In particular, we will use that the Dirichlet trace operator can be extended by continuity to surjective mappings

$$\gamma_D : H_{\Delta}^{3/2}(\Omega) \rightarrow H^1(\partial\Omega) \quad \text{and} \quad \gamma_D : H_{\Delta}^1(\Omega) \rightarrow H^{1/2}(\partial\Omega), \quad (4.4)$$

and the Neumann trace operator can be extended by continuity to surjective mappings

$$\gamma_N : H_{\Delta}^{3/2}(\Omega) \rightarrow L^2(\partial\Omega) \quad \text{and} \quad \gamma_N : H_{\Delta}^1(\Omega) \rightarrow H^{-1/2}(\partial\Omega); \quad (4.5)$$

cf. [62, Theorems 7.3 and 7.4, Chapter 2] for the case of a bounded smooth domain and, e.g. [49, Lemma 3.1 and Lemma 3.2]. At the same time the second Green's identity

$$(-\Delta f, g)_{L^2(\Omega)} - (f, -\Delta g)_{L^2(\Omega)} = (\gamma_D f, \gamma_N g)_{L^2(\partial\Omega)} - (\gamma_N f, \gamma_D g)_{L^2(\partial\Omega)}, \quad (4.6)$$

well known for $f, g \in H^2(\Omega)$, remains valid for $f, g \in H_{\Delta}^{3/2}(\Omega)$ and extends further to functions $f, g \in H_{\Delta}^1(\Omega)$

$$(-\Delta f, g)_{L^2(\Omega)} - (f, -\Delta g)_{L^2(\Omega)} = \langle \gamma_D f, \gamma_N g \rangle - \langle \gamma_N f, \gamma_D g \rangle, \quad (4.7)$$

where $\langle \cdot, \cdot \rangle$ denotes the extension of the $L^2(\partial\Omega)$ -inner product onto the dual pair $H^{1/2}(\partial\Omega) \times H^{-1/2}(\partial\Omega)$ and $H^{-1/2}(\partial\Omega) \times H^{1/2}(\partial\Omega)$, respectively. As usual, here

$$H^{1/2}(\partial\Omega) \hookrightarrow L^2(\partial\Omega) \hookrightarrow H^{-1/2}(\partial\Omega) \quad (4.8)$$

is viewed as a rigging of Hilbert spaces, that is, some uniformly positive self-adjoint operator j in $L^2(\partial\Omega)$ with $\text{dom}(j) = H^{1/2}(\partial\Omega)$ is fixed and viewed as an isomorphism

$$j : H^{1/2}(\partial\Omega) \longrightarrow L^2(\partial\Omega). \quad (4.9)$$

As scalar product on $H^{1/2}(\partial\Omega)$ we choose $(\varphi, \psi)_{H^{1/2}(\partial\Omega)} := (j\varphi, j\psi)_{L^2(\partial\Omega)}$; it follows that $H^{-1/2}(\partial\Omega)$ coincides with the completion of $L^2(\partial\Omega)$ with respect to $(j^{-1}\cdot, j^{-1}\cdot)_{L^2(\partial\Omega)}$, and j^{-1} admits an extension to an isomorphism

$$\widetilde{j^{-1}} : H^{-1/2}(\partial\Omega) \longrightarrow L^2(\partial\Omega).$$

The inner product $\langle \cdot, \cdot \rangle$ on the right hand side of (4.7) is

$$\langle \varphi, \psi \rangle := (j\varphi, \widetilde{j^{-1}\psi})_{L^2(\partial\Omega)}, \quad \varphi \in H^{1/2}(\partial\Omega), \quad \psi \in H^{-1/2}(\partial\Omega), \quad (4.10)$$

and extends the $L^2(\partial\Omega)$ scalar product in the sense that $\langle \varphi, \psi \rangle = (\varphi, \psi)_{L^2(\partial\Omega)}$ for $\varphi \in H^{1/2}(\partial\Omega)$ and $\psi \in L^2(\partial\Omega)$. A standard and convenient choice for j in (4.9) in many situations is

$$j_\Delta := (-\Delta_{\partial\Omega} + I)^{1/4} : H^{1/2}(\partial\Omega) \longrightarrow L^2(\partial\Omega), \quad (4.11)$$

where $-\Delta_{\partial\Omega}$ denotes the Laplace–Beltrami operator in $L^2(\partial\Omega)$. In this case

$$\widetilde{j_\Delta^{-1}} = (-\Delta_{\partial\Omega} + I)^{-1/4} : H^{-1/2}(\partial\Omega) \longrightarrow L^2(\partial\Omega);$$

cf. Remark 4.5 for other natural choices of j . Note in this connection that j_Δ maps $H^s(\partial\Omega)$ isomorphically onto $H^{s-1/2}(\partial\Omega)$ for any $s \in \mathbb{R}$.

In this context we also recall the following lemma, which is essentially a consequence of the asymptotics of the eigenvalues of the Laplace–Beltrami operator on compact manifolds; cf. [4, Proof of Proposition 5.4.1], [5, Theorem 2.1.2], and [17, Lemma 4.7].

Lemma 4.1. *Let \mathcal{K} be a Hilbert space and assume that $X \in \mathcal{B}(\mathcal{K}, H^s(\partial\Omega))$ has the property $\text{ran } X \subset H^r(\partial\Omega)$ for some $r > s \geq 0$. Then*

$$X \in \mathcal{S}_{\frac{n-1}{r-s}}(\mathcal{K}, H^s(\partial\Omega))$$

and hence $X \in \mathfrak{S}_p(\mathcal{K}, H^s(\partial\Omega))$ for $p > \frac{n-1}{r-s}$.

As a useful consequence of Lemma 4.1 we note that for $r > 0$ the canonical embeddings $\iota_r : H^r(\partial\Omega) \longrightarrow L^2(\partial\Omega)$ and $\iota_{-r} : L^2(\partial\Omega) \longrightarrow H^{-r}(\partial\Omega)$ satisfy

$$\iota_r \in \mathcal{S}_{\frac{n-1}{r}}(H^r(\partial\Omega), L^2(\partial\Omega)) \quad \text{and} \quad \iota_{-r} \in \mathcal{S}_{\frac{n-1}{r}}(L^2(\partial\Omega), H^{-r}(\partial\Omega)),$$

respectively. In fact, the assertion for the embedding ι_r follows after fixing a unitary operator $U : L^2(\partial\Omega) \longrightarrow H^r(\partial\Omega)$, applying Lemma 4.1 to the operator $X = \iota_r U$ and noting that the singular values of X and ι_r are the same. Since the dual operator $\iota'_r : L^2(\partial\Omega) \longrightarrow H^{-r}(\partial\Omega)$ coincides with the canonical embedding ι_{-r} of $L^2(\partial\Omega)$ into $H^{-r}(\partial\Omega)$ the second assertion follows. By composition and (1.4) we also conclude

$$\iota_{-r} \circ \iota_r \in \mathcal{S}_{\frac{n-1}{2r}}(H^r(\partial\Omega), H^{-r}(\partial\Omega)). \quad (4.12)$$

4.2. Schrödinger operators with Dirichlet, Neumann, and Robin boundary conditions

Let $\Omega \subset \mathbb{R}^n$ be an exterior domain as in Section 4.1. In the following we consider a Schrödinger differential expression with a bounded, measurable, real valued potential V ,

$$\mathcal{L} = -\Delta + V, \quad V \in L^\infty(\Omega). \quad (4.13)$$

With the expression in (4.13) one naturally associates the minimal operator

$$\begin{aligned} S_{\min} f &= \mathcal{L} f, \\ \operatorname{dom}(S_{\min}) &= H_0^2(\Omega) = \{f \in H^2(\Omega) : \gamma_D f = \gamma_N f = 0\}, \end{aligned} \quad (4.14)$$

and the maximal operator

$$\begin{aligned} S_{\max} f &= \mathcal{L} f, \\ \operatorname{dom}(S_{\max}) &= \{f \in L^2(\Omega) : -\Delta f + V f \in L^2(\Omega)\}, \end{aligned}$$

in $L^2(\Omega)$; the expression Δf in $\operatorname{dom}(S_{\max})$ is understood in the sense of distributions. We note that $\operatorname{dom}(S_{\max})$ equipped with the graph norm coincides with the Hilbert space $H_\Delta^0(\Omega)$ introduced above. In the next lemma we collect some well-known properties of S_{\min} and S_{\max} ; for the simplicity of S we refer to [22, Proposition 2.2] and we mention that another proof of this fact can be obtained following the reasoning in [28, Example 5.3]. The density of $H_\Delta^s(\Omega)$ in $\operatorname{dom}(S^*)$ equipped with the graph norm is shown (for the case of a bounded domain) in [62, Chapter 2, Theorem 6.4].

Lemma 4.2. *The operator $S := S_{\min}$ is a densely defined, closed, simple, symmetric operator in $L^2(\Omega)$. The deficiency indices of S coincide and are both infinite,*

$$\dim(\operatorname{ran}(S - i)^\perp) = \dim(\operatorname{ran}(S + i)^\perp) = \infty.$$

The adjoint of the minimal operator is the maximal operator,

$$S^* = S_{\min}^* = S_{\max} \quad \text{and} \quad S = S_{\min} = S_{\max}^*,$$

and the spaces $H_\Delta^s(\Omega)$, $s \in [0, 2]$, are dense in $\operatorname{dom}(S^)$ equipped with the graph norm.*

In Sections 4.3 and 4.4 we are interested in scattering systems consisting of different self-adjoint realizations of \mathcal{L} in $L^2(\Omega)$. The self-adjoint Dirichlet and Neumann operators associated to the densely defined, semibounded, closed quadratic forms

$$\begin{aligned} \mathfrak{a}_D[f, g] &= (\nabla f, \nabla g)_{(L^2(\Omega))^n} + (Vf, g)_{L^2(\Omega)}, & \operatorname{dom}(\mathfrak{a}_D) &= H_0^1(\Omega), \\ \mathfrak{a}_N[f, g] &= (\nabla f, \nabla g)_{(L^2(\Omega))^n} + (Vf, g)_{L^2(\Omega)}, & \operatorname{dom}(\mathfrak{a}_N) &= H^1(\Omega), \end{aligned}$$

are given by

$$\begin{aligned} A_D f &= \mathcal{L} f, & \operatorname{dom}(A_D) &= \{f \in H^2(\Omega) : \gamma_D f = 0\}, \\ A_N f &= \mathcal{L} f, & \operatorname{dom}(A_N) &= \{f \in H^2(\Omega) : \gamma_N f = 0\}, \end{aligned} \quad (4.15)$$

and for a real valued function $\alpha \in L^\infty(\partial\Omega)$ the quadratic form

$$\mathfrak{a}_\alpha[f, g] = \mathfrak{a}_N[f, g] - (\alpha\gamma_D f, \gamma_D g)_{L^2(\partial\Omega)}, \quad \text{dom}(\mathfrak{a}_\alpha) = H^1(\Omega),$$

is also densely defined, closed and semibounded from below, and hence gives rise to a semibounded self-adjoint operator in $L^2(\Omega)$, which has the form

$$A_\alpha f = \mathcal{L}f, \quad \text{dom}(A_\alpha) = \{f \in H_\Delta^{3/2}(\Omega) : \alpha\gamma_D f = \gamma_N f\}. \quad (4.16)$$

We remark that the H^2 -regularity of the functions in $\text{dom}(A_D)$ and $\text{dom}(A_N)$ is a classical fact (see the monographs [3, 61, 62]) and the $H^{3/2}$ -regularity of the functions in $\text{dom}(A_\alpha)$ can be found in, e.g. [14, Corollary 6.25]; in the case that the coefficient α in the Robin boundary condition is continuously differentiable also $\text{dom}(A_\alpha)$ is contained in $H^2(\Omega)$; cf. [68, Theorem 4.18].

4.3. Scattering matrix for the Dirichlet and Robin realization

In this subsection we consider the pair $\{A_D, A_\alpha\}$ consisting of the self-adjoint Dirichlet and Robin operator associated to \mathcal{L} in (4.15) and (4.16) on an exterior domain $\Omega \subset \mathbb{R}^2$; here we restrict ourselves to the two dimensional situation in order to ensure that the trace class condition (3.1) for the resolvent difference is satisfied; cf. Remark 4.4.

Before formulating and proving our main result on the system $\{A_D, A_\alpha\}$ we recall the definition and some useful properties of the Dirichlet-to-Neumann map. First we note that for any $\psi \in H^{1/2}(\partial\Omega)$ and $z \in \rho(A_D)$ there exists a unique solution $f_z \in H_\Delta^1(\Omega)$ of the boundary value problem

$$-\Delta f_z + V f_z = z f_z, \quad \gamma_D f_z = \psi \in H^{1/2}(\partial\Omega). \quad (4.17)$$

The corresponding solution operator is given by

$$P_D(z) : H^{1/2}(\partial\Omega) \longrightarrow H_\Delta^1(\Omega) \subset L^2(\Omega), \quad \psi \mapsto f_z. \quad (4.18)$$

For $z \in \rho(A_D)$ the *Dirichlet-to-Neumann map* $\Lambda_{1/2}(z)$ is defined by

$$\Lambda_{1/2}(z) : H^{1/2}(\partial\Omega) \longrightarrow H^{-1/2}(\partial\Omega), \quad \psi \mapsto \gamma_N P_D(z)\psi, \quad (4.19)$$

and takes Dirichlet boundary values $\gamma_D f_z$ of the solution $f_z \in H_\Delta^1(\Omega)$ of (4.17) to their Neumann boundary values $\gamma_N f_z \in H^{-1/2}(\partial\Omega)$.

Now we are ready to formulate and prove a representation of the scattering matrix for the pair $\{A_D, A_\alpha\}$.

Theorem 4.3. *Let $\Omega \subset \mathbb{R}^2$ be an exterior domain with a C^∞ -smooth boundary, let $V \in L^\infty(\Omega)$ and $\alpha \in L^\infty(\partial\Omega)$ be real valued functions, and let A_D and A_α be the*

self-adjoint Dirichlet and Robin realizations of $\mathcal{L} = -\Delta + V$ in $L^2(\Omega)$ in (4.15) and (4.16), respectively. Moreover, let $\Lambda_{1/2}(\cdot)$ be the Dirichlet-to-Neumann map defined in (4.19) and let

$$M_\alpha^D(z) := \widetilde{j^{-1}(\alpha - \Lambda_{1/2}(z))}j^{-1}, \quad z \in \rho(A_D), \quad (4.20)$$

where $j : H^{1/2}(\partial\Omega) \rightarrow L^2(\partial\Omega)$ denotes some uniformly positive self-adjoint operator in $L^2(\partial\Omega)$ with $\text{dom}(j) = H^{1/2}(\partial\Omega)$ as in (4.8)–(4.9).

Then $\{A_D, A_\alpha\}$ is a complete scattering system and

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda), \quad \mathcal{H}_\lambda := \overline{\text{ran}(\text{Im } M_\alpha^D(\lambda + i0))},$$

forms a spectral representation of A_D^{ac} such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A_D, A_\alpha; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A_D, A_\alpha\}$ admits the representation

$$S(A_D, A_\alpha; \lambda) = I_{\mathcal{H}_\lambda} - 2i\sqrt{\text{Im } M_\alpha^D(\lambda + i0)} M_\alpha^D(\lambda + i0)^{-1} \sqrt{\text{Im } M_\alpha^D(\lambda + i0)}.$$

Proof. It follows from (4.15) and (4.16) that the operator $A_\alpha \cap A_D$ coincides with the minimal operator $S = S_{\min}$ associated with \mathcal{L} in (4.14), which is closed, densely defined and simple by Lemma 4.2. Define the operator T as a restriction of S^* to the domain $H_\Delta^1(\Omega)$,

$$Tf = -\Delta f + Vf, \quad \text{dom}(T) = H_\Delta^1(\Omega),$$

and let

$$\Gamma_0 f := j\gamma_D f \quad \text{and} \quad \Gamma_1 f := \widetilde{j^{-1}(\alpha\gamma_D - \gamma_N)}f, \quad f \in \text{dom}(T). \quad (4.21)$$

We claim that $\Pi_\alpha^D = \{L^2(\partial\Omega), \Gamma_0, \Gamma_1\}$ is a B -generalized boundary triple for S^* with the \mathfrak{S}_1 -regular Weyl function $M_\alpha^D(\cdot)$ given by (4.20) such that

$$A_D = T \upharpoonright \ker(\Gamma_0) \quad \text{and} \quad A_\alpha = T \upharpoonright \ker(\Gamma_1). \quad (4.22)$$

In fact, for $f, g \in \text{dom}(T)$ we use (4.7) and the fact that α is real valued, and compute

$$\begin{aligned} & (\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g) \\ &= (\widetilde{j^{-1}(\alpha\gamma_D - \gamma_N)}f, j\gamma_D g) - (j\gamma_D f, \widetilde{j^{-1}(\alpha\gamma_D - \gamma_N)}g) \\ &= \langle \alpha\gamma_D f - \gamma_N f, \gamma_D g \rangle - \langle \gamma_D f, \alpha\gamma_D g - \gamma_N g \rangle \\ &= \langle \gamma_D f, \gamma_N g \rangle - \langle \gamma_N f, \gamma_D g \rangle \\ &= (Tf, g) - (f, Tg) \end{aligned}$$

and hence Green's identity (2.1) is satisfied. Moreover, $\gamma_D : \text{dom}(T) \rightarrow H^{1/2}(\partial\Omega)$ is well defined and surjective according to (4.4), and since $j : H^{1/2}(\partial\Omega) \rightarrow L^2(\partial\Omega)$ is an

isomorphism we conclude $\text{ran}(\Gamma_0) = L^2(\partial\Omega)$, i.e., Γ_0 is surjective. From Lemma 4.2 we directly obtain that $\text{dom}(T) = H^1_\Delta(\Omega)$ is dense in $\text{dom}(S^*)$ equipped with the graph norm (which is equal to the space $H^0_\Delta(\Omega)$) and hence we have $\overline{T} = S^*$. Moreover, it follows from Green's identity (2.1) that the restrictions $T \upharpoonright \ker(\Gamma_0)$ and $T \upharpoonright \ker(\Gamma_1)$ are both symmetric operators in $L^2(\Omega)$ and from the definition of the boundary maps it is clear that the self-adjoint operators A_D and A_α are contained in the symmetric operators $T \upharpoonright \ker(\Gamma_0)$ and $T \upharpoonright \ker(\Gamma_1)$, and hence they coincide. Therefore, $\Pi^D_\alpha = \{L^2(\partial\Omega), \Gamma_0, \Gamma_1\}$ is a B -generalized boundary triple for S^* such that (4.22) holds.

In order to see that the Weyl function is given by (4.20) recall that $\Lambda_{1/2}(z)\gamma_D f_z = \gamma_N f_z$ for $f_z \in \ker(T - z)$, $z \in \rho(A_D)$, according to the definition of the Dirichlet-to-Neumann map $\Lambda_{1/2}(\cdot)$ in (4.19). Hence we obtain

$$\widetilde{J^{-1}}(\alpha - \Lambda_{1/2}(z))J^{-1}\Gamma_0 f_z = \widetilde{J^{-1}}(\alpha\gamma_D f_z - \Lambda_{1/2}(z)\gamma_D f_z) = \Gamma_1 f_z$$

for $f_z \in \ker(T - z)$ and $z \in \rho(A_D)$, and this yields (4.20).

It remains to verify that $M^D_\alpha(\cdot) \in \mathfrak{S}_1$ -regular. For this we denote the γ -field associated to Π^D_α by $\gamma^D_\alpha(\cdot)$ and use the relation (2.7) with some $\xi \in \rho(A_D) \cap \rho(A_\alpha) \cap \rho(A_N) \cap \mathbb{R}$ and all $z \in \rho(A_D)$. Observe that (2.6), $\xi = \bar{\xi}$, and the choice of Γ_1 in (4.21) yield

$$\gamma^D_\alpha(\xi)^* h = \Gamma_1(A_D - \xi)^{-1} h = -\widetilde{J^{-1}}\gamma_N(A_D - \xi)^{-1} h \quad (4.23)$$

for all $h \in L^2(\Omega)$. Since $\text{dom}(A_D) \subset H^2(\Omega)$ we conclude from (4.1) that the range of the mapping $\gamma_N(A_D - \xi)^{-1}$ is contained in $H^{1/2}(\partial\Omega)$. As γ_N maps $H^2(\Omega)$ continuously onto $H^{1/2}(\partial\Omega)$ (cf. (4.1)) this operator is defined on the whole space $L^2(\Omega)$ and

$$\gamma_N(A_D - \xi)^{-1} \in \mathcal{B}(L^2(\Omega), H^{1/2}(\partial\Omega)).$$

Now we use that the canonical embedding operator $\iota_{-1/2} \circ \iota_{1/2} : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega)$ is compact and belongs to $\mathcal{S}_1(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))$ by (4.12). Thus we have

$$\gamma_N(A_D - \xi)^{-1} \in \mathcal{S}_1(L^2(\Omega), H^{-1/2}(\partial\Omega))$$

and hence (4.23) yields $\gamma^D_\alpha(\xi)^* \in \mathcal{S}_1(L^2(\Omega), L^2(\partial\Omega))$. It follows that also $\gamma^D_\alpha(\xi) \in \mathcal{S}_1(L^2(\partial\Omega), L^2(\Omega))$ and hence by (2.5) for all $z \in \rho(A_D)$

$$\gamma^D_\alpha(z) = (I + (z - \xi)(A_D - z)^{-1})\gamma^D_\alpha(\xi) \in \mathcal{S}_1(L^2(\partial\Omega), L^2(\Omega)). \quad (4.24)$$

Therefore

$$(z - \xi)\gamma^D_\alpha(\xi)^*\gamma^D_\alpha(z) \in \mathcal{S}_{1/2}(L^2(\partial\Omega)), \quad z \in \rho(A_D). \quad (4.25)$$

Since $\mathcal{S}_{1/2}(L^2(\partial\Omega)) \subset \mathfrak{S}_1(L^2(\partial\Omega))$ and $M^D_\alpha(\xi) = M^D_\alpha(\xi)^*$ we conclude from (2.7) and (4.25) that

$$K(z) := M_\alpha^D(z) - M_\alpha^D(\xi) = (z - \xi)\gamma_\alpha^D(\xi)^* \gamma_\alpha^D(z) \in \mathfrak{S}_1(L^2(\partial\Omega)), \quad z \in \mathbb{C}_+.$$

Since $M_\alpha^D(\cdot)$ is a strict Nevanlinna function, $K(\cdot)$ is a strict Nevanlinna function too. It remains to show that

$$C := M_\alpha^D(\xi) = \widetilde{j^{-1}\alpha j^{-1}} - \widetilde{j^{-1}\Lambda_{1/2}(\xi)j^{-1}}$$

is boundedly invertible. Using that the maps (4.4) and (4.5) are surjective and $\xi \in \rho(A_D) \cap \rho(A_N) \cap \mathbb{R}$ we find that the self-adjoint operator $\widetilde{j^{-1}\Lambda_{1/2}(\xi)j^{-1}}$ is surjective, and hence boundedly invertible in $L^2(\partial\Omega)$. From $\text{ran}(\alpha j^{-1}) \subseteq L^2(\partial\Omega)$ we obtain that $\widetilde{j^{-1}\alpha j^{-1}}$ is compact and therefore $M_\alpha^D(\xi)$ is a Fredholm operator. Furthermore, $\ker(M_\alpha^D(\xi)) \neq \{0\}$ by Lemma 2.14 and hence $C = M_\alpha^D(\xi)$ is boundedly invertible. Therefore $M_\alpha^D(\cdot)$ is an \mathfrak{S}_1 -regular Weyl function. Now the assertions in Theorem 4.3 follow from Theorem 3.1. \square

Remark 4.4. For $n = 2, 3, 4, \dots$ one obtains in the same way as in the proof of Theorem 4.3 using (4.12) that

$$\gamma_\alpha^D(z) \in \mathcal{S}_{n-1}(L^2(\partial\Omega), L^2(\Omega)) \quad \text{and} \quad \gamma_\alpha^D(z)^* \in \mathcal{S}_{n-1}(L^2(\Omega), L^2(\partial\Omega))$$

for all $z \in \rho(A_D)$ and since $M_\alpha^D(z)^{-1} \in \mathcal{B}(L^2(\partial\Omega))$, $z \in \rho(A_D) \cap \rho(A_\alpha)$, we conclude from Krein's formula in Proposition 2.7 (iii) that

$$(A_\alpha - z)^{-1} - (A_D - z)^{-1} = -\gamma_\alpha^D(z)M_\alpha^D(z)^{-1}\gamma_\alpha^D(\bar{z})^* \in \mathcal{S}_{\frac{n-1}{2}}(L^2(\Omega)) \quad (4.26)$$

for all $z \in \rho(A_D) \cap \rho(A_\alpha)$ by Proposition 2.7 (iii). In particular, for $n = 2$ one gets the \mathfrak{S}_1 -resolvent comparability of A_α and A_D . This well known result goes back to Birman [24] (see also [17,45,53,54,63] for more details on singular value estimates in this context).

Remark 4.5. There are several possibilities to choose the operator j in (4.9) used for the extension (4.10) of the $L^2(\partial\Omega)$ scalar product in the rigging (4.8). Besides the choice $j_\Delta = (-\Delta_{\partial\Omega} + I)^{1/4}$ in (4.11) the following choice is very convenient for the scattering matrix, since it allows to express it completely in terms of the Dirichlet-to-Neumann map: Fix some $\lambda_0 < \min\{\sigma(A_D), \sigma(A_N)\}$ and note that the restriction $\Lambda_1(\lambda_0)$ (see also the beginning of Section 5.4) of the Dirichlet-to-Neumann map $\Lambda_{1/2}(\lambda_0)$ onto $H^1(\partial\Omega)$ is a non-negative self-adjoint operator in $L^2(\partial\Omega)$ with a bounded everywhere defined inverse $\Lambda_1(\lambda_0)^{-1}$ in $L^2(\partial\Omega)$; the Neumann-to-Dirichlet map. Then also the square root $\sqrt{\Lambda_1(\lambda_0)}$ is a non-negative self-adjoint operator in $L^2(\partial\Omega)$ which is boundedly invertible, and we have $\text{dom}(\sqrt{\Lambda_1(\lambda_0)}) = H^{1/2}(\partial\Omega)$ (see, e.g., [18, Proposition 3.2 (iii)]). Hence

$$j = \sqrt{\Lambda_1(\lambda_0)} : H^{1/2}(\partial\Omega) \longrightarrow L^2(\partial\Omega)$$

is a possible choice for the definition of the scalar product $\langle \cdot, \cdot \rangle$ in (4.10).

Following [23, Section 1] one defines the adjoint X^+ of an operator

$$X \in \mathcal{B}(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))$$

in the rigging $H^{1/2}(\partial\Omega) \hookrightarrow L^2(\partial\Omega) \hookrightarrow H^{-1/2}(\partial\Omega)$ via

$$\langle X\varphi, \psi \rangle = \langle \varphi, X^+\psi \rangle, \quad \varphi, \psi \in H^{1/2}(\partial\Omega).$$

The imaginary part of the operator X is defined by $\operatorname{Im} X = \frac{1}{2i}(X - X^+)$. The operator X is self-adjoint if $X = X^+$ and X is non-negative if $\langle X\varphi, \varphi \rangle \geq 0$ for all $\varphi \in H^{1/2}(\partial\Omega)$.

From the fact that the function $M_\alpha^D(\cdot)$ in (4.20) is \mathfrak{S}_1 -regular with values in $\mathcal{B}(L^2(\partial\Omega))$ we conclude

$$\Lambda_{1/2}(z) \in \mathcal{B}(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega)), \quad z \in \mathbb{C}_+.$$

Together with Lemma 2.6 this yields the following corollary.

Corollary 4.6. *Let $\Omega \subset \mathbb{R}^2$ be an exterior domain with a C^∞ -smooth boundary and let $\Lambda_{1/2}(\cdot)$ be the Dirichlet-to-Neumann map defined in (4.19). Then the following holds.*

- (i) *The limit $\Lambda_{1/2}(\lambda + i0) = \lim_{\varepsilon \rightarrow +0} \Lambda_{1/2}(\lambda + i\varepsilon)$ exists for a.e. $\lambda \in \mathbb{R}$ in the norm of $\mathcal{B}(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))$;*
- (ii) *$\Lambda_{1/2}(\lambda + i0) \in \mathcal{B}(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))$ is boundedly invertible for a.e. $\lambda \in \mathbb{R}$;*
- (iii) *$\Lambda_{1/2}(\lambda + i\varepsilon) - \Lambda_{1/2}(\lambda + i0) \in \mathfrak{S}_p(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))$ for $p \in (1, \infty]$, $\varepsilon > 0$ and a.e. $\lambda \in \mathbb{R}$, and*

$$\lim_{\varepsilon \rightarrow +0} \|\Lambda_{1/2}(\lambda + i\varepsilon) - \Lambda_{1/2}(\lambda + i0)\|_{\mathfrak{S}_p(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))} = 0;$$

- (iv) *$\operatorname{Im} \Lambda_{1/2}(\lambda + i0) = \lim_{\varepsilon \rightarrow +0} \operatorname{Im} \Lambda_{1/2}(\lambda + i\varepsilon)$ exists for a.e. $\lambda \in \mathbb{R}$ in the $\mathfrak{S}_1(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))$ -norm and $-\operatorname{Im} \Lambda_{1/2}(\lambda + i0) \geq 0$.*

4.4. Scattering matrix for the Neumann and Robin realization

In this subsection we discuss a representation of the scattering matrix for the pair $\{A_N, A_\alpha\}$ consisting of the self-adjoint Neumann and Robin operator associated to \mathcal{L} in (4.15) and (4.16). Here Ω is an exterior domain in \mathbb{R}^2 or \mathbb{R}^3 ; it is known from [24] (for \mathbb{R}^2) and [15, 58] (for \mathbb{R}^2 and \mathbb{R}^3) that the trace class condition (3.1) for the resolvent difference is satisfied; cf. Remark 4.8.

In a similar way as in the previous subsection we first define the Neumann-to-Dirichlet map $\mathcal{N}(z)$ as an operator in $L^2(\partial\Omega)$ for all $z \in \rho(A_N)$. Recall first that for $\varphi \in L^2(\partial\Omega)$ and $z \in \rho(A_N)$ the boundary value problem

$$-\Delta f_z + V f_z = z f_z, \quad \gamma_N f_z = \varphi, \tag{4.27}$$

admits a unique solution $f_z \in H_{\Delta}^{3/2}(\Omega)$. The corresponding solution operator is given by

$$P_N(z) : L^2(\partial\Omega) \longrightarrow H_{\Delta}^{3/2}(\Omega) \subset L^2(\Omega), \quad \varphi \mapsto f_z. \quad (4.28)$$

For $z \in \rho(A_N)$ the *Neumann-to-Dirichlet map* is defined by

$$\mathcal{N}(z) : L^2(\partial\Omega) \longrightarrow L^2(\partial\Omega), \quad \varphi \mapsto \gamma_D P_N(z)\varphi. \quad (4.29)$$

It is clear that $\mathcal{N}(z)$ maps Neumann boundary values $\gamma_N f_z$ of the solutions $f_z \in H_{\Delta}^{3/2}(\Omega)$ of (4.27) onto their Dirichlet boundary values $\gamma_D f_z$; here γ_N and γ_D denote the extensions of the Dirichlet and Neumann trace operators onto $H_{\Delta}^{3/2}(\Omega)$ from (4.4) and (4.5), respectively. Since (4.27) admits a unique solution for each $\varphi \in L^2(\partial\Omega)$ it is clear that the operators $P_N(z)$ and $\mathcal{N}(z)$ are well defined on $L^2(\partial\Omega)$.

In the next theorem the scattering matrix of the pair $\{A_N, A_{\alpha}\}$ is expressed in terms of the limit values of the Neumann-to-Dirichlet map $\mathcal{N}(z)$ and the parameter α in the boundary condition of the Robin realization A_{α} . In contrast to Theorem 4.3 here it is also assumed that $\alpha^{-1} \in L^{\infty}(\partial\Omega)$.

Theorem 4.7. *Let $\Omega \subset \mathbb{R}^n$, $n = 2, 3$, be an exterior domain with a C^{∞} -smooth boundary, let $V \in L^{\infty}(\Omega)$ and $\alpha \in L^{\infty}(\partial\Omega)$ be real valued functions such that $\alpha^{-1} \in L^{\infty}(\partial\Omega)$, and let A_N and A_{α} be the self-adjoint Neumann and Robin realizations of $\mathcal{L} = -\Delta + V$ in $L^2(\Omega)$ in (4.15) and (4.16), respectively. Moreover, let $\mathcal{N}(\cdot)$ be the Neumann-to-Dirichlet map defined in (4.29).*

Then $\{A_N, A_{\alpha}\}$ is a complete scattering system and

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_{\lambda}), \quad \mathcal{H}_{\lambda} := \overline{\text{ran}(\text{Im } \mathcal{N}(\lambda + i0))},$$

forms a spectral representation of A_N^{ac} such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A_N, A_{\alpha}; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A_N, A_{\alpha}\}$ admits the representation

$$S(A_N, A_{\alpha}; \lambda) = I_{\mathcal{H}_{\lambda}} + 2i\sqrt{\text{Im } \mathcal{N}(\lambda + i0)} (I - \alpha \mathcal{N}(\lambda + i0))^{-1} \alpha \sqrt{\text{Im } \mathcal{N}(\lambda + i0)}.$$

Proof. First we note that the assumption $\alpha^{-1} \in L^{\infty}(\partial\Omega)$ implies $A_N \cap A_{\alpha} = S$, where S is the minimal operator associated to \mathcal{L} in (4.14). Recall that S is closed, densely defined and simple by Lemma 4.2. Define the operator T as a restriction of S^* by

$$Tf = -\Delta f + Vf, \quad \text{dom}(T) = H_{\Delta}^{3/2}(\Omega),$$

and let

$$\Gamma_0 f := \gamma_N f \quad \text{and} \quad \Gamma_1 f := \gamma_D f - \frac{1}{\alpha} \gamma_N f, \quad f \in \text{dom}(T). \quad (4.30)$$

We claim that $\Pi_{\alpha}^N = \{L^2(\partial\Omega), \Gamma_0, \Gamma_1\}$ is a B -generalized boundary triple for S^* with the \mathfrak{S}_1 -regular Weyl function

$$M_\alpha^N(z) = \mathcal{N}(z) - \frac{1}{\alpha}, \quad z \in \rho(A_N), \quad (4.31)$$

such that

$$A_N = T \upharpoonright \ker(\Gamma_0) \quad \text{and} \quad A_\alpha = T \upharpoonright \ker(\Gamma_1). \quad (4.32)$$

In fact, Green's identity (2.1) is an immediate consequence of the definition of the boundary mappings and (4.6), and $\text{ran } \Gamma_0 = L^2(\partial\Omega)$ holds by (4.5). Moreover, $\text{dom}(T)$ is dense in $\text{dom}(S^*)$ with respect to the graph norm by Lemma 4.2 and $A_\alpha = T \upharpoonright \ker(\Gamma_1)$ is clear from (4.16). Furthermore, the self-adjoint operator A_N in (4.15) is contained in $T \upharpoonright \ker(\Gamma_0)$ and since the latter is symmetric (a consequence of Green's identity (2.1)) both operators coincide, that is, (4.32) holds, and Π_α^N is a B -generalized boundary triple. For $f_z \in \ker(T - z)$, $z \in \rho(A_N)$, we have

$$\left(\mathcal{N}(z) - \frac{1}{\alpha}\right) \Gamma_0 f_z = \mathcal{N}(z) \gamma_N f_z - \frac{1}{\alpha} \gamma_N f_z = \gamma_D f_z - \frac{1}{\alpha} \gamma_N f_z = \Gamma_1 f_z$$

and hence the Weyl function $M_\alpha^N(\cdot)$ corresponding to Π_α^N is given by (4.31).

It remains to check that the Weyl function $M_\alpha^N(\cdot)$ is \mathfrak{S}_1 -regular. This is done in a similar way as in Theorem 4.3. Denote the γ -field associated to Π_α^N by $\gamma_\alpha^N(\cdot)$ and use (cf. (2.7))

$$M_\alpha^N(z) = M_\alpha^N(\xi)^* + (z - \bar{\xi}) \gamma_\alpha^N(\xi)^* \gamma_\alpha^N(z) \quad (4.33)$$

with some fixed $\xi \in \rho(A_N) \cap \rho(A_\alpha) \cap \mathbb{R}$ and all $z \in \rho(A_N)$. From (4.30), (4.15), and (4.1) we obtain for any $f \in L^2(\Omega)$

$$\gamma_\alpha^N(\xi)^* f = \Gamma_1(A_N - \xi)^{-1} f = \gamma_D(A_N - \xi)^{-1} f \in H^{3/2}(\partial\Omega)$$

and hence Lemma 4.1 yields

$$\gamma_\alpha^N(\xi)^* \in \mathcal{S}_{\frac{2(n-1)}{3}}(L^2(\Omega), L^2(\partial\Omega)) \quad (4.34)$$

and

$$\gamma_\alpha^N(z) \in \mathcal{S}_{\frac{2(n-1)}{3}}(L^2(\partial\Omega), L^2(\Omega)) \quad (4.35)$$

for all $z \in \rho(A_N)$. Now combining (1.4) with (4.33) yields

$$K(z) := M_\alpha^N(z) - M_\alpha^N(\xi) = (z - \xi) \gamma_\alpha^N(\xi)^* \gamma_\alpha^N(z) \in \mathcal{S}_{\frac{n-1}{3}}(L^2(\partial\Omega))$$

for $z \in \rho(A_N)$. Since $\mathcal{S}_{(n-1)/3}(L^2(\partial\Omega))$ is contained in $\mathfrak{S}_1(L^2(\partial\Omega))$ for $n = 2, 3$, and $M_\alpha^N(\xi) = M_\alpha^N(\xi)^*$ we conclude that $K(z) \in \mathfrak{S}_1(L^2(\partial\Omega))$, $z \in \mathbb{C}_+$. Because $M_\alpha^N(\cdot)$ is a strict Nevanlinna function $K(\cdot)$ is also strict. Let us show that

$$C := M_\alpha^N(\xi) = \mathcal{N}(\xi) - \frac{1}{\alpha}$$

is invertible. In fact, since $\frac{1}{\alpha}$ is a boundedly invertible operator and $\mathcal{N}(\xi)$ is a compact operator it follows that $M_\alpha^N(\xi)$ is a Fredholm operator. Furthermore, $\ker(M_\alpha^N(\xi))$ is trivial by Lemma 2.14 and hence C is boundedly invertible. Therefore, the Weyl function $M_\alpha^N(\cdot)$ is \mathfrak{S}_1 -regular. Now the assertions in Theorem 4.7 follow from Theorem 3.1,

$$\operatorname{Im} M_\alpha^N(z) = \operatorname{Im} \mathcal{N}(z), \quad M_\alpha^N(z)^{-1} = -(I - \alpha \mathcal{N}(z))^{-1} \alpha, \quad z \in \mathbb{C}_+,$$

and

$$\operatorname{Im} M_\alpha^N(\lambda + i0) = \operatorname{Im} \mathcal{N}(\lambda + i0), \quad M_\alpha^N(\lambda + i0)^{-1} = -(I - \alpha \mathcal{N}(\lambda + i0))^{-1} \alpha$$

for a.e. $\lambda \in \mathbb{R}$. \square

Remark 4.8. From (4.34) and (4.35) one concludes in the same way as in Remark 4.4 that Krein's formula in Proposition 2.7 (iii) and the property (1.4) leads to

$$(A_\alpha - z)^{-1} - (A_N - z)^{-1} = -\gamma_\alpha^N(z) M_\alpha^N(z)^{-1} \gamma_\alpha^N(\bar{z})^* \in \mathcal{S}_{\frac{n-1}{3}}(L^2(\Omega)) \quad (4.36)$$

for all $z \in \rho(A_\alpha) \cap \rho(A_N)$; cf. [15,58]. Note that a weaker estimate with $\mathcal{S}_{\frac{n-1}{2}}$ instead of $\mathcal{S}_{\frac{n-1}{3}}$ is immediate from (4.26) first established by Birman [24] (see Remark 4.4). It yields the \mathfrak{S}_1 -resolvent comparability for $n = 2$.

Remark 4.9. The definition of the boundary triples Π_α^D and Π_α^N in Theorems 4.3 and 4.7 given for an exterior domain Ω , and the form and properties of the corresponding Weyl functions remain the same in the case of a bounded domain Ω with smooth boundary. The constructions and properties are only based on the compactness and smoothness of $\partial\Omega$.

5. Schrödinger operators with interactions supported on hypersurfaces

In this section we investigate scattering systems consisting of Schrödinger operators in \mathbb{R}^n . Here the Euclidean space is decomposed into a smooth bounded domain and its complement, and the usual self-adjoint Schrödinger operator on the whole space is compared with the orthogonal sum of the Dirichlet or Neumann operators on the subdomains in Section 5.2 and 5.3, and with a Schrödinger operator with a singular δ -potential supported on the interface in Section 5.4. In our main results Theorem 5.1, 5.4, and 5.6 we obtain explicit forms of the scattering matrices in terms of Dirichlet-to-Neumann or Neumann-to-Dirichlet maps. As in Section 4 the strategy in the proofs is to apply the general result Theorem 3.1 to suitable B -generalized boundary triples. Here we shall assume for convenience that a simplicity condition for the underlying symmetric operator is satisfied; this condition can be dropped in which case Corollary 3.3 would yield

a slightly more involved representation of the scattering matrix. We also refer the interested reader to [Remarks 5.2, 5.5, and 5.7](#), where singular value estimates due to Birman, Grubb and others are revisited.

5.1. Preliminaries on orthogonal sums and couplings of Schrödinger operators

Let $\Omega_- \subset \mathbb{R}^n$ be a bounded domain with C^∞ -smooth boundary $\partial\Omega_-$ and let $\Omega_+ := \mathbb{R}^n \setminus \overline{\Omega_-}$ be the corresponding C^∞ -smooth exterior domain. Denote the common boundary of Ω_+ and Ω_- by $\mathcal{C} := \partial\Omega_\pm$. Throughout this section we consider a Schrödinger differential expression with a bounded, measurable, real valued potential V on \mathbb{R}^n ,

$$\mathcal{L} = -\Delta + V, \quad V \in L^\infty(\mathbb{R}^n). \quad (5.1)$$

In the following we shall adapt the notation from [Section 4.1](#) in an obvious way, e.g. $H^s(\Omega_\pm)$ and $H^r(\mathcal{C})$ denote the Sobolev spaces on Ω_\pm and the common boundary (or interface) \mathcal{C} , respectively, the spaces $H_\Delta^s(\Omega_\pm)$, $s \in [0, 2]$, are defined and equipped with scalar products as in [\(4.2\)–\(4.3\)](#), and we shall use the notation

$$H_\Delta^s(\mathbb{R}^n \setminus \mathcal{C}) := H_\Delta^s(\Omega_+) \times H_\Delta^s(\Omega_-), \quad s \in [0, 2].$$

A function $f : \mathbb{R}^n \rightarrow \mathbb{C}$ is often written in a two component form $f = \{f^+, f^-\}$, where $f^\pm : \Omega_\pm \rightarrow \mathbb{C}$ denote the restrictions of f onto Ω_\pm . The Dirichlet and Neumann trace operators will be denoted by γ_D^\pm and γ_N^\pm , and we emphasize that the Neumann trace is taken with respect to the outer normal of Ω_\pm . In particular, $\gamma_N^+ f^+ + \gamma_N^- f^- = 0$ for a function $f = \{f^+, f^-\} \in H^2(\mathbb{R}^n)$. We also note that the mapping properties of the Dirichlet and Neumann trace operators in [\(4.4\)](#) and [\(4.5\)](#) are valid for both domains Ω_+ and Ω_- , and the same is true for the extensions of Green's identity in [\(4.6\)](#) and [\(4.7\)](#), respectively. Furthermore, we shall use in the proofs in [Section 5.2](#) and [Section 5.3](#) that γ_D^\pm and γ_N^\pm admit continuous extensions

$$\gamma_D^\pm : H_\Delta^0(\Omega_\pm) \rightarrow H^{-1/2}(\mathcal{C}) \quad \text{and} \quad \gamma_N^\pm : H_\Delta^0(\Omega_\pm) \rightarrow H^{-3/2}(\mathcal{C})$$

and that Green's identity extends to $f_\pm \in H^2(\Omega_\pm)$ and $g_\pm \in H_\Delta^0(\Omega_\pm)$ in the form

$$(-\Delta f_\pm, g_\pm)_{L^2(\Omega_\pm)} - (f_\pm, -\Delta g_\pm)_{L^2(\Omega_\pm)} = \langle \gamma_D^\pm f_\pm, \gamma_N^\pm g_\pm \rangle - \langle \gamma_N^\pm f_\pm, \gamma_D^\pm g_\pm \rangle; \quad (5.2)$$

cf. [\[62\]](#) and [\[52, Chapter I, Theorem 3.3 and Corollary 3.3\]](#). In [\(5.2\)](#) the inner products $\langle \cdot, \cdot \rangle$ on the right hand side denote the continuations of the $L^2(\mathcal{C})$ inner product onto $H^{3/2}(\mathcal{C}) \times H^{-3/2}(\mathcal{C})$ and $H^{1/2}(\mathcal{C}) \times H^{-1/2}(\mathcal{C})$, respectively, and in the following it will always be clear from the context which duality is used; cf. [\(4.8\)–\(4.10\)](#).

The differential expression [\(5.1\)](#) induces self-adjoint operators in $L^2(\mathbb{R}^n)$. The natural self-adjoint realization is the free Schrödinger operator,

$$A_{\text{free}}f = \mathcal{L}f, \quad \text{dom}(A_{\text{free}}) = H^2(\mathbb{R}^n), \quad (5.3)$$

which is semibounded from below. Clearly the functions in $\text{dom}(A_{\text{free}})$ do not reflect the decomposition of \mathbb{R}^n into the domains Ω_+ and Ω_- . Furthermore, we will make use of the self-adjoint orthogonal sum

$$A_D = A_D^+ \oplus A_D^-, \quad \text{dom}(A_D) = \{f = \{f^+, f^-\} \in H^2(\Omega_+) \oplus H^2(\Omega_-) : \gamma_D^+ f^+ = \gamma_D^- f^- = 0\}, \quad (5.4)$$

of the self-adjoint Dirichlet operators A_D^\pm in $L^2(\Omega_\pm)$ in (4.15), and of the self-adjoint orthogonal sum

$$A_N = A_N^+ \oplus A_N^-, \quad \text{dom}(A_N) = \{f = \{f^+, f^-\} \in H^2(\Omega_+) \oplus H^2(\Omega_-) : \gamma_N^+ f^+ = \gamma_N^- f^- = 0\}, \quad (5.5)$$

of the self-adjoint Neumann operators A_N^\pm in $L^2(\Omega_\pm)$ in (4.15). We shall sometimes refer to A_D as Dirichlet realization of \mathcal{L} with respect to \mathcal{C} and to A_N as Neumann realization of \mathcal{L} with respect to \mathcal{C} . The properties of A_D^\pm and A_N^\pm extend in a natural way to their orthogonal sums A_D and A_N in (5.4) and (5.5), respectively. In particular, the Dirichlet realization A_D and the Neumann realization A_N of \mathcal{L} with respect to \mathcal{C} are both semibounded from below.

5.2. Scattering matrix for the free Schrödinger operator and the Dirichlet realization with respect to \mathcal{C}

We shall derive a representation for the scattering matrix of the scattering system $\{A_D, A_{\text{free}}\}$ in \mathbb{R}^2 . Let $\Lambda_{1/2}^\pm(z) : H^{1/2}(\mathcal{C}) \mapsto H^{-1/2}(\mathcal{C})$ be the Dirichlet-to-Neumann map defined in (4.19) with respect to Ω_\pm , that is,

$$\Lambda_{1/2}^\pm(z) \gamma_D^\pm f_z^\pm = \gamma_N^\pm f_z^\pm \quad (5.6)$$

holds for any solution $f_z^\pm \in H^1(\Omega_\pm)$ of the equation $-\Delta f_z^\pm + V_\pm f_z^\pm = z f_z^\pm$ and $z \in \rho(A_D^\pm)$. Furthermore, define the operator-valued function $\Lambda_{1/2}(\cdot)$ by

$$\Lambda_{1/2}(z) := \Lambda_{1/2}^+(z) + \Lambda_{1/2}^-(z) : H^{1/2}(\mathcal{C}) \longrightarrow H^{-1/2}(\mathcal{C}), \quad z \in \rho(A_D). \quad (5.7)$$

Theorem 5.1. *Let $\Omega_\pm \subset \mathbb{R}^2$ be as above, let $V \in L^\infty(\mathbb{R}^2)$ be a real valued function, and let A_{free} and A_D be the self-adjoint Schrödinger operators in $L^2(\mathbb{R}^2)$ in (5.3) and (5.4), respectively. Moreover, let $\Lambda_{1/2}(\cdot)$ be given by (5.7) and let*

$$M_{\text{free}}^D(z) := -\widetilde{j^{-1}} \Lambda_{1/2}(z) j^{-1}, \quad z \in \mathbb{C}_+, \quad (5.8)$$

where $j : H^{1/2}(\mathcal{C}) \rightarrow L^2(\mathcal{C})$ denotes some uniformly positive self-adjoint operator in $L^2(\mathcal{C})$ with $\text{dom}(j) = H^{1/2}(\mathcal{C})$ as in (4.8)–(4.9).

Then $\{A_D, A_{\text{free}}\}$ is a complete scattering system. If the densely defined, closed, symmetric operator $S := A_D \cap A_{\text{free}}$ has no eigenvalues then

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda), \quad \mathcal{H}_\lambda := \overline{\text{ran}(\text{Im } M_{\text{free}}^D(\lambda + i0))},$$

forms a spectral representation of A_D^{ac} such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A_D, A_{\text{free}}; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A_D, A_{\text{free}}\}$ admits the representation

$$S(A_D, A_{\text{free}}; \lambda) = I_{\mathcal{H}_\lambda} - 2i\sqrt{\text{Im } M_{\text{free}}^D(\lambda + i0)}M_{\text{free}}^D(\lambda + i0)^{-1}\sqrt{\text{Im } M_{\text{free}}^D(\lambda + i0)}.$$

Proof. The closed symmetric operator $S = A_D \cap A_{\text{free}}$ in $L^2(\mathbb{R}^2)$ is given by

$$\begin{aligned} Sf &= \mathcal{L}f, \\ \text{dom}(S) &= \{f = \{f^+, f^-\} \in H^2(\mathbb{R}^2) : \gamma_D^+ f^+ = \gamma_D^- f^- = 0\}. \end{aligned} \quad (5.9)$$

It is clear that S is a closed extension of the orthogonal sum of the minimal operators $S^+ \oplus S^-$ associated to the restriction of \mathcal{L} onto Ω_+ and Ω_- as in (4.14) and Lemma 4.2. It follows that S is densely defined and since we have assumed that S has no eigenvalues it follows from [21, Corollary 4.4] that S is simple. We claim that the adjoint S^* is given by

$$\begin{aligned} S^*f &= \mathcal{L}f, \\ \text{dom}(S^*) &= \{f = \{f^+, f^-\} \in H_\Delta^0(\mathbb{R}^2 \setminus \mathcal{C}) : \gamma_D^+ f^+ = \gamma_D^- f^-\}. \end{aligned}$$

In fact, since $S^* \subset (S^+)^* \oplus (S^-)^*$ it follows that

$$\text{dom}(S^*) \subset H_\Delta^0(\mathbb{R}^2 \setminus \mathcal{C}) = \text{dom}(S^+)^* \times \text{dom}(S^-)^*$$

and that $S^*f = \mathcal{L}f$ for $f \in \text{dom}(S^*)$. Therefore, we only have to verify that $f = \{f^+, f^-\} \in \text{dom}(S^*)$ satisfies the interface condition

$$\gamma_D^+ f^+ = \gamma_D^- f^-. \quad (5.10)$$

Assume that for $f = \{f^+, f^-\} \in \text{dom}(S^*)$ and all $h = \{h^+, h^-\} \in \text{dom}(S)$ we have

$$(Sh, f)_{L^2(\mathbb{R}^2)} = (h, S^*f)_{L^2(\mathbb{R}^2)},$$

that is,

$$\begin{aligned} (-\Delta h^+, f^+)_{L^2(\Omega_+)} + (-\Delta h^-, f^-)_{L^2(\Omega_-)} \\ = (h^+, -\Delta f^+)_{L^2(\Omega_+)} + (h^-, -\Delta f^-)_{L^2(\Omega_-)}. \end{aligned}$$

Then it follows from Green's identity (5.2) and the conditions

$$\gamma_D^\pm h^\pm = 0 \quad \text{and} \quad \gamma_N^+ h^+ + \gamma_N^- h^- = 0$$

that

$$\begin{aligned} 0 &= (-\Delta h^+, f^+)_{L^2(\Omega_+)} - (h^+, -\Delta f^+)_{L^2(\Omega_+)} \\ &\quad + (-\Delta h^-, f^-)_{L^2(\Omega_-)} - (h^-, -\Delta f^-)_{L^2(\Omega_-)} \\ &= \langle \gamma_D^+ h^+, \gamma_N^+ f^+ \rangle - \langle \gamma_N^+ h^+, \gamma_D^+ f^+ \rangle + \langle \gamma_D^- h^-, \gamma_N^- f^- \rangle - \langle \gamma_N^- h^-, \gamma_D^- f^- \rangle \\ &= \langle \gamma_N^- h^-, \gamma_D^+ f^+ - \gamma_D^- f^- \rangle \end{aligned}$$

holds for all $h = \{h^+, h^-\} \in \text{dom}(S)$. This implies (5.10).

Now we proceed in a similar manner as in the proofs of Theorem 4.3 and Theorem 4.7 in the previous section. We consider the operator T defined as a restriction of S^* by

$$\begin{aligned} Tf &= \mathcal{L}f, \\ \text{dom}(T) &= \{f = \{f^+, f^-\} \in H_\Delta^1(\mathbb{R}^2 \setminus \mathcal{C}) : \gamma_D^+ f^+ = \gamma_D^- f^-\}, \end{aligned}$$

and for $f \in \text{dom}(T)$ we agree on the notation

$$\gamma_D f := \gamma_D^+ f^+ = \gamma_D^- f^-, \quad f = \{f^+, f^-\} \in \text{dom}(T). \quad (5.11)$$

We claim that $\Pi_{\text{free}}^D = \{L^2(\mathcal{C}), \Gamma_0, \Gamma_1\}$, where

$$\Gamma_0 f := J \gamma_D f \quad \text{and} \quad \Gamma_1 f := -\widetilde{J^{-1}} (\gamma_N^+ f^+ + \gamma_N^- f^-), \quad f \in \text{dom}(T),$$

is a B -generalized boundary triple with an \mathfrak{S}_1 -regular Weyl function given by (5.8) such that

$$A_D = T \upharpoonright \ker(\Gamma_0) \quad \text{and} \quad A_{\text{free}} = T \upharpoonright \ker(\Gamma_1). \quad (5.12)$$

In fact, for $f = \{f^+, f^-\}$, $g = \{g^+, g^-\} \in \text{dom}(T)$ we compute with the help of Green's identity (4.7) and (4.10) that

$$\begin{aligned} &(\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g) \\ &= \langle -\gamma_N^+ f^+ - \gamma_N^- f^-, \gamma_D g \rangle - \langle \gamma_D f, -\gamma_N^+ g^+ - \gamma_N^- g^- \rangle \\ &= \langle \gamma_D^+ f^+, \gamma_N^+ g^+ \rangle - \langle \gamma_N^+ f^+, \gamma_D^+ g^+ \rangle + \langle \gamma_D^- f^-, \gamma_N^- g^- \rangle - \langle \gamma_N^- f^-, \gamma_D^- g^- \rangle \\ &= (-\Delta f^+, g^+) - (f^+, -\Delta g^+) + (-\Delta f^-, g^-) - (f^-, -\Delta g^-) \\ &= (Tf, g) - (f, Tg) \end{aligned}$$

and (4.4) implies $\text{ran}(\Gamma_0) = L^2(\mathcal{C})$ in the present situation; cf. the proof of Theorem 4.3. Since $T \upharpoonright \ker(\Gamma_0)$ and $T \upharpoonright \ker(\Gamma_1)$ are both symmetric operators by (2.1), and contain the self-adjoint operators A_D and A_{free} , respectively, it follows that (5.12) is satisfied. Furthermore, as $S = A_D \cap A_{\text{free}}$ it is clear that the self-adjoint operators A_D and A_{free} are disjoint extensions of S . It follows from Proposition 2.9 that

$$(\text{dom}(A_D) + \text{dom}(A_{\text{free}})) \subset \text{dom}(T) \quad (5.13)$$

is dense in $\text{dom}(S^*)$ with respect to the graph norm. Hence $\overline{T} = S^*$. Therefore Π_{free}^D is B -generalized boundary triple such that (5.12) holds.

Next we show that the Weyl function $M_{\text{free}}^D(\cdot)$ corresponding to Π_{free}^D is \mathfrak{S}_1 -regular and has the form (5.8). Let $f_z = \{f_z^+, f_z^-\} \in \ker(T - z)$, $z \in \rho(A_D)$, and use (5.6) and (5.7) to compute

$$\begin{aligned} -\widetilde{j^{-1}}\Lambda_{1/2}(z)j^{-1}\Gamma_0 f_z &= -\widetilde{j^{-1}}(\Lambda_{1/2}^+(z) + \Lambda_{1/2}(z)^-)\gamma_D f_z \\ &= -\widetilde{j^{-1}}(\gamma_N^+ f_z^+ + \gamma_N^- f_z^-) = \Gamma_1 f_z. \end{aligned}$$

Hence the Weyl function is $M_{\text{free}}^D(z) = -\widetilde{j^{-1}}\Lambda_{1/2}(z)j^{-1}$. In order to see that $M_{\text{free}}^D(\cdot)$ is \mathfrak{S}_1 -regular we proceed in the same way as in the proof of Theorem 4.3. Let $\gamma_{\text{free}}^D(\cdot)$ be the γ -field corresponding to the B -generalized boundary triple Π_{free}^D and use

$$M_{\text{free}}^D(z) = M_{\text{free}}^D(\xi)^* + (z - \xi)\gamma_{\text{free}}^D(\xi)^*\gamma_{\text{free}}^D(z) \quad (5.14)$$

(see (2.7)) with some $\xi \in \rho(A_D) \cap \rho(A_{\text{free}}) \cap (-\infty, \text{ess inf } V)$ and all $z \in \rho(A_D)$. For $h = \{h^+, h^-\} \in L^2(\mathbb{R}^n)$ we have

$$\begin{aligned} \gamma_{\text{free}}^D(\xi)^* h &= \Gamma_1(A_D - \xi)^{-1} h \\ &= -\widetilde{j^{-1}}(\gamma_N^+(A_D^+ - \xi)^{-1} h^+ + \gamma_N^-(A_D^- - \xi)^{-1} h^-) \end{aligned} \quad (5.15)$$

and since $\text{dom}(A_D) \subset H^2(\Omega_+) \times H^2(\Omega_-)$ we conclude from (4.1) that

$$\gamma_N^+(A_D^+ - \xi)^{-1} h^+ + \gamma_N^-(A_D^- - \xi)^{-1} h^- \in H^{1/2}(\mathcal{C}).$$

As in the proof of Theorem 4.3 it then follows from (5.15) and (4.12) with $r = 1/2$ and $n = 2$ that

$$\gamma_{\text{free}}^D(\xi)^* \in \mathcal{S}_1(L^2(\mathbb{R}^2), L^2(\mathcal{C})) \quad (5.16)$$

and $\gamma_{\text{free}}^D(z) \in \mathcal{S}_1(L^2(\mathcal{C}), L^2(\mathbb{R}^2))$ for all $z \in \rho(A_D)$. Hence (5.14) yields that

$$K(z) := M_{\text{free}}^D(z) - M_{\text{free}}^D(\xi) \in \mathfrak{S}_1(L^2(\mathcal{C})), \quad z \in \mathbb{C}_+,$$

where it was used that $M_{\text{free}}^D(\xi)^* = M_{\text{free}}^D(\xi)$. It follows that $M_{\text{free}}^D(\xi)$ is a Fredholm operator since $0 \in \rho(M_{\text{free}}^D(z))$ for $z \in \mathbb{C}_+$. On the other hand by [Lemma 2.14](#) we have $\ker(M_{\text{free}}^D(\xi)) = \{0\}$ for $\xi \in \rho(A_D) \cap \rho(A_{\text{free}}) \cap (-\infty, \text{ess inf } V)$. Thus $M_{\text{free}}^D(\xi)$ is boundedly invertible which shows that $M_{\text{free}}^D(\cdot)$ is \mathfrak{S}_1 -regular. Now the assertions follow directly from [Theorem 3.1](#). \square

Remark 5.2. As in [Remarks 4.4 and 4.8](#) it follows from [\(5.15\)](#) and [\(4.12\)](#) in the same way as in [\(5.16\)](#) that for $n \geq 2$

$$\gamma_{\text{free}}^D(z)^* \in \mathcal{S}_{n-1}(L^2(\mathbb{R}^n), L^2(\mathcal{C}))$$

for $z \in \rho(A_D)$. This yields $\gamma_{\text{free}}^D(z) \in \mathcal{S}_{n-1}(L^2(\mathcal{C}), L^2(\mathbb{R}^n))$ for $z \in \rho(A_D)$ and hence Krein's formula in [Proposition 2.7](#) (iii) implies

$$(A_{\text{free}} - z)^{-1} - (A_D - z)^{-1} = -\gamma_{\text{free}}^D(z) M_{\text{free}}^D(z)^{-1} \gamma_{\text{free}}^D(\bar{z})^* \in \mathcal{S}_{\frac{n-1}{2}}(L^2(\mathbb{R}^n))$$

for all $z \in \rho(A_{\text{free}}) \cap \rho(A_D)$; cf. [\[24,54\]](#). For further development with applications to the scattering theory we also refer the reader to [\[33\]](#) and [\[79\]](#).

Remark 5.3. As in [Remark 4.5](#) there is a particularly convenient choice of the operator j in [\(4.8\)–\(4.9\)](#) in the present context. Namely, since for any $z < \min\{\sigma(A_D^\pm), \sigma(A_N^\pm)\}$ the self-adjoint operators

$$\sqrt{\Lambda_{1/2}^+(z)} \quad \text{and} \quad \sqrt{\Lambda_{1/2}^-(z)}$$

defined on $H^{1/2}(\mathcal{C})$ are non-negative and boundedly invertible in $L^2(\mathcal{C})$ it follows that

$$j := \sqrt{\Lambda_{1/2}^+(z)} + \sqrt{\Lambda_{1/2}^-(z)} : H^{1/2}(\mathcal{C}) \longrightarrow L^2(\mathcal{C})$$

is a possible choice for the definition of the inner product $\langle \cdot, \cdot \rangle$ in [\(4.10\)](#).

5.3. Scattering matrix for the free Schrödinger operator and the Neumann realization with respect to \mathcal{C}

In this section we consider the pair $\{A_N, A_{\text{free}}\}$ consisting of the orthogonal sum $A_N = A_N^+ \oplus A_N^-$ of the Neumann operators in [\(5.5\)](#) and the free Schrödinger operator in [\(5.3\)](#). We first define the Neumann-to-Dirichlet maps

$$\mathcal{N}_{-1/2}^\pm(z) : H^{-1/2}(\mathcal{C}) \longrightarrow H^{1/2}(\mathcal{C}), \quad z \in \rho(A_N),$$

as extensions of the Neumann-to-Dirichlet maps on $L^2(\mathcal{C})$ defined in the beginning of [Section 4.4](#). More precisely, we recall that for $\phi^\pm \in H^{-1/2}(\mathcal{C})$ and $z \in \rho(A_N^\pm)$ the boundary value problem

$$-\Delta f^\pm + V_\pm f^\pm = z f^\pm, \quad \gamma_N^\pm f^\pm = \phi^\pm, \quad (5.17)$$

admits a unique solution $f_z^\pm \in H_\Delta^1(\Omega_\pm)$. The corresponding solution operator is denoted by

$$\mathcal{P}_N^\pm(z) : H^{-1/2}(\mathcal{C}) \longrightarrow H_\Delta^1(\mathcal{C}) \subset L^2(\mathcal{C}), \quad \phi^\pm \mapsto f_z^\pm.$$

Note that the restriction of $\mathcal{P}_N^\pm(z)$ onto $L^2(\mathcal{C})$ coincides with the solution operator defined in (4.28). For $z \in \rho(A_N^\pm)$ the *Neumann-to-Dirichlet map* is defined by

$$\mathcal{N}_{-1/2}^\pm(z) : H^{-1/2}(\mathcal{C}) \longrightarrow H^{1/2}(\mathcal{C}), \quad \phi^\pm \mapsto \gamma_D^\pm \mathcal{P}_N^\pm(z) \phi^\pm. \quad (5.18)$$

Clearly, $\mathcal{N}_{-1/2}^\pm(z)$ is an extension of the Neumann-to-Dirichlet map defined in (4.29) onto $H^{-1/2}(\mathcal{C})$, the operators in (5.18) map Neumann boundary values $\gamma_N^\pm f_z^\pm$ of solutions $f_z^\pm \in H_\Delta^1(\Omega_\pm)$ of (5.17) to the corresponding Dirichlet boundary values $\gamma_D^\pm f_z^\pm \in H^{1/2}(\mathcal{C})$.

In the next theorem we obtain an expression for the scattering matrix of the pair $\{A_N, A_{\text{free}}\}$ in terms of the sum

$$\mathcal{N}_{-1/2}(z) := \mathcal{N}_{-1/2}^+(z) + \mathcal{N}_{-1/2}^-(z) : H^{-1/2}(\mathcal{C}) \longrightarrow H^{1/2}(\mathcal{C}), \quad z \in \rho(A_N), \quad (5.19)$$

of the Neumann-to-Dirichlet maps in (5.18).

Theorem 5.4. *Let $\Omega_\pm \subset \mathbb{R}^2$ be as above, let $V \in L^\infty(\mathbb{R}^2)$ be a real valued function, and let A_{free} and A_N be the self-adjoint Schrödinger operators in $L^2(\mathbb{R}^2)$ in (5.3) and (5.5), respectively. Moreover, let $\mathcal{N}_{-1/2}(\cdot)$ be given by (5.19) and let*

$$M_{\text{free}}^N(z) := j \mathcal{N}_{-1/2}(z) \widetilde{j}, \quad z \in \mathbb{C}_+, \quad (5.20)$$

where $j : H^{1/2}(\mathcal{C}) \longrightarrow L^2(\mathcal{C})$ denotes some uniformly positive self-adjoint operator in $L^2(\mathcal{C})$ with $\text{dom}(j) = H^{1/2}(\mathcal{C})$ as in (4.8)–(4.9).

Then $\{A_N, A_{\text{free}}\}$ is a complete scattering system. If the densely defined, closed, symmetric operator $S := A_N \cap A_{\text{free}}$ has no eigenvalues then

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda), \quad \mathcal{H}_\lambda := \overline{\text{ran}(\text{Im } M_{\text{free}}^N(\lambda + i0))},$$

forms a spectral representation of A_N^{ac} such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A_N, A_{\text{free}}; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A_N, A_{\text{free}}\}$ admits the representation

$$S(A_N, A_{\text{free}}; \lambda) = I_{\mathcal{H}_\lambda} - 2i \sqrt{\text{Im } M_{\text{free}}^N(\lambda + i0)} M_{\text{free}}^N(\lambda + i0)^{-1} \sqrt{\text{Im } M_{\text{free}}^N(\lambda + i0)}.$$

Proof. The proof of Theorem 5.4 is very similar to the proof of Theorem 5.1, and hence we present a sketch only. Consider the closed symmetric operator $S = A_N \cap A_{\text{free}}$ in $L^2(\mathbb{R}^2)$ which is given by

$$Sf = \mathcal{L}f,$$

$$\operatorname{dom}(S) = \{f = \{f^+, f^-\} \in H^2(\mathbb{R}^2) : \gamma_N^+ f^+ = \gamma_N^- f^- = 0\}.$$

It follows that S is densely defined, the assumption $\sigma_p(S) = \emptyset$ and same arguments as in [21, Proof of Lemma 4.3] ensure that S is simple, and a similar consideration as in the proof of Theorem 5.1 shows that the adjoint S^* is given by

$$S^*f = \mathcal{L}f,$$

$$\operatorname{dom}(S^*) = \{f = \{f^+, f^-\} \in H_\Delta^0(\mathbb{R}^2 \setminus \mathcal{C}) : \gamma_N^+ f^+ = \gamma_N^- f^-\}.$$

Next we consider the operator T defined as a restriction of S^* by

$$Tf = \mathcal{L}f,$$

$$\operatorname{dom}(T) = \{f = \{f^+, f^-\} \in H_\Delta^1(\mathbb{R}^2 \setminus \mathcal{C}) : \gamma_N^+ f^+ = \gamma_N^- f^-\}.$$

As in the proof of Theorem 5.1 one verifies that $\Pi_{\text{free}}^N = \{L^2(\mathcal{C}), \Gamma_0, \Gamma_1\}$, where

$$\Gamma_0 f := \widetilde{j^{-1} \gamma_N^+ f^+} \quad \text{and} \quad \Gamma_1 f := j(\gamma_D^+ f^+ - \gamma_D^- f^-), \quad f \in \operatorname{dom}(T),$$

is a B -generalized boundary triple with the Weyl function $M_{\text{free}}^N(\cdot)$ given by (5.20) such that

$$A_N = T \upharpoonright \ker(\Gamma_0) \quad \text{and} \quad A_{\text{free}} = T \upharpoonright \ker(\Gamma_1).$$

Let us show that the Weyl function $M_{\text{free}}^N(\cdot)$ is \mathfrak{S}_1 -regular. Denote the γ -field corresponding to the B -generalized boundary triple Π_{free}^N by $\gamma_{\text{free}}^N(\cdot)$ and use

$$M_{\text{free}}^N(z) = M_{\text{free}}^N(\xi)^* + (z - \xi)\gamma_{\text{free}}^N(\xi)^*\gamma_{\text{free}}^N(z) \quad (5.21)$$

with some fixed $\xi \in \rho(A_N) \cap \rho(A_{\text{free}}) \cap (-\infty, \operatorname{ess\,inf} V)$ and all $z \in \rho(A_N)$. From (4.1) and $\operatorname{dom}(A_N) \subset H^2(\Omega_+) \times H^2(\Omega_-)$ we conclude for $h = \{h^+, h^-\} \in L^2(\mathbb{R}^n)$ that

$$j^{-1}\gamma_{\text{free}}^N(\xi)^*h = j^{-1}\Gamma_1(A_N - \xi)^{-1}h$$

$$= \gamma_D^+(A_N^+ - \xi)^{-1}h^+ - \gamma_D^-(A_N^- - \xi)^{-1}h^- \in H^{3/2}(\mathcal{C}). \quad (5.22)$$

Since $j^{-1}\gamma_{\text{free}}^N(\xi)^* \in \mathcal{B}(L^2(\mathbb{R}^2), H^{1/2}(\mathcal{C}))$, Lemma 4.1 applies with $r = 3/2$, $s = 1/2$ and gives

$$j^{-1}\gamma_{\text{free}}^N(\xi)^* \in \mathcal{S}_1(L^2(\mathbb{R}^2), H^{1/2}(\mathcal{C}))$$

and since j is an isomorphism from $H^{1/2}(\mathcal{C})$ onto $L^2(\mathcal{C})$,

$$\gamma_{\text{free}}^N(\xi)^* \in \mathcal{S}_1(L^2(\mathbb{R}^2), L^2(\mathcal{C})). \quad (5.23)$$

Therefore $\gamma_{\text{free}}^N(z) \in \mathcal{S}_1(L^2(\mathcal{C}), L^2(\mathbb{R}^2))$ for all $z \in \rho(A_N)$. Now it follows from (5.21) that

$$K(z) := M_{\text{free}}^N(z) - M_{\text{free}}^N(\xi) \in \mathcal{S}_{1/2}(L^2(\mathcal{C})) \subset \mathfrak{S}_1(L^2(\mathcal{C})), \quad z \in \mathbb{C}_+,$$

where we have used that $M_{\text{free}}^N(\xi) = M_{\text{free}}^N(\xi)^*$. It remains to show that $M_{\text{free}}^N(\xi)$ is invertible, which follows from the same reasoning as in the end of the proof of Theorem 5.1. Hence $M_{\text{free}}^N(\cdot)$ is \mathfrak{S}_1 -regular and the assertions of Theorem 5.4 follow directly from Theorem 3.1. \square

Remark 5.5. As in Remark 5.2 the considerations in (5.22) and (5.23) together with Lemma 4.1 show that for $n \geq 2$

$$\gamma_{\text{free}}^N(z)^* \in \mathcal{S}_{n-1}(L^2(\mathbb{R}^n), L^2(\mathcal{C})), \quad \gamma_{\text{free}}^N(z) \in \mathcal{S}_{n-1}(L^2(\mathcal{C}), L^2(\mathbb{R}^n))$$

for all $z \in \rho(A_N)$. Hence

$$(A_{\text{free}} - z)^{-1} - (A_N - z)^{-1} = -\gamma_{\text{free}}^N(z) M_{\text{free}}^N(z)^{-1} \gamma_{\text{free}}^N(\bar{z})^* \in \mathcal{S}_{\frac{n-1}{2}}(L^2(\mathbb{R}^n))$$

for all $z \in \rho(A_{\text{free}}) \cap \rho(A_N)$. The latter gives another proof of a result of Grubb from [54].

5.4. Schrödinger operators with δ -potentials supported on hypersurfaces

In this third and last application on scattering matrices for coupled Schrödinger operators we consider the pair $\{A_{\text{free}}, A_{\delta, \alpha}\}$, where $\alpha \in L^\infty(\mathcal{C})$ is a real valued function and $A_{\delta, \alpha}$ is a Schrödinger operator with δ -potential of strength α supported on the hypersurface \mathcal{C} defined by

$$A_{\delta, \alpha} f = -\Delta f + V f, \\ \text{dom}(A_{\delta, \alpha}) = \left\{ f = \begin{pmatrix} f^+ \\ f^- \end{pmatrix} \in H_{\Delta}^{3/2}(\mathbb{R}^n \setminus \mathcal{C}) : \begin{array}{l} \gamma_D^+ f^+ = \gamma_D^- f^-, \\ \alpha \gamma_D^\pm f^\pm = \gamma_N^+ f^+ + \gamma_N^- f^- \end{array} \right\}. \quad (5.24)$$

Such type of Schrödinger operators with singular interactions have attracted a lot of attention in the past; cf. [39] for a survey and e.g. [16] for further references and an approach via boundary mappings closely related to the present considerations. According to [16, Theorem 3.5, Proposition 3.7, and Theorem 3.16] the operator $A_{\delta, \alpha}$ in (5.24) is self-adjoint in $L^2(\mathbb{R}^n)$, semibounded from below and coincides with the self-adjoint operator associated to the closed sesquilinear form

$$\mathfrak{a}_{\delta, \alpha}[f, g] = (\nabla f, \nabla g) + (V f, g) - (\alpha \gamma_D^\pm f, \gamma_D^\pm g)_{L^2(\mathcal{C})}, \quad f, g \in H^1(\mathbb{R}^n).$$

We define the Dirichlet-to-Neumann maps

$$\Lambda_1^\pm(z) : H^1(\mathcal{C}) \longrightarrow L^2(\mathcal{C}), \quad z \in \rho(A_D^\pm),$$

as restrictions of the Dirichlet-to-Neumann maps on $H^{1/2}(\mathcal{C})$ in (4.19); cf. Remark 4.5. More precisely, for $\phi^\pm \in H^1(\mathcal{C})$ and $z \in \rho(A_D^\pm)$ the boundary value problem

$$-\Delta f^\pm + V_\pm f^\pm = z f^\pm, \quad \gamma_D^\pm f^\pm = \phi^\pm,$$

admits a unique solution $f_z^\pm \in H_\Delta^{3/2}(\Omega_\pm)$. The corresponding solution operators are denoted by

$$\mathcal{P}_D^\pm(z) : H^1(\mathcal{C}) \longrightarrow H_\Delta^{3/2}(\mathcal{C}) \subset L^2(\mathcal{C}), \quad \phi^\pm \mapsto f_z^\pm,$$

and it is clear that the restriction of $\mathcal{P}_D^\pm(z)$ in (4.18) onto $H^1(\mathcal{C})$ coincides with $\mathcal{P}_D^\pm(z)$. For $z \in \rho(A_D^\pm)$ the Dirichlet-to-Neumann maps $\Lambda_1^\pm(\cdot)$ on $H^1(\mathcal{C})$ are given by

$$\Lambda_1^\pm(z) : H^1(\mathcal{C}) \longrightarrow L^2(\mathcal{C}), \quad \phi^\pm \mapsto \gamma_N^\pm \mathcal{P}_D^\pm(z) \phi^\pm, \quad (5.25)$$

and by construction $\Lambda_1^\pm(z)$ are the restrictions of the Dirichlet-to-Neumann maps $\Lambda_{1/2}^\pm(z)$ in (4.19) onto $H^1(\mathcal{C})$.

In the next theorem we obtain an expression for the scattering matrix of the pair $\{A_{\text{free}}, A_{\delta, \alpha}\}$ in terms of the sum

$$\Lambda_1(z) := \Lambda_1^+(z) + \Lambda_1^-(z) : H^1(\mathcal{C}) \longrightarrow L^2(\mathcal{C}), \quad z \in \rho(A_D), \quad (5.26)$$

of the Dirichlet-to-Neumann maps in (5.25). Theorem 5.6 and its proof can be viewed as a variant of Theorem 4.7; in the same way as in Theorem 4.7 it is assumed that $\alpha^{-1} \in L^\infty(\mathcal{C})$.

Theorem 5.6. *Let $\Omega_\pm \subset \mathbb{R}^n$, $n = 2, 3$, be as above, let $V \in L^\infty(\mathbb{R}^n)$ and $\alpha \in L^\infty(\mathcal{C})$ be real valued functions such that $\alpha^{-1} \in L^\infty(\mathcal{C})$, and let A_{free} and $A_{\delta, \alpha}$ be the self-adjoint realizations of the Schrödinger expression given by (5.3) and (5.24), respectively. Moreover, let $\Lambda_1(\cdot)$ be as in (5.26).*

Then $\{A_{\text{free}}, A_{\delta, \alpha}\}$ is a complete scattering system. If the densely defined, closed, symmetric operator $S := A_{\text{free}} \cap A_{\delta, \alpha}$ has no eigenvalues then

$$L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda), \quad \mathcal{H}_\lambda := \overline{\text{ran}(\text{Im}(\Lambda_1(\lambda + i0))^{-1})},$$

forms a spectral representation of $A_{\text{free}}^{\text{ac}}$ such that for a.e. $\lambda \in \mathbb{R}$ the scattering matrix $\{S(A_{\text{free}}, A_{\delta, \alpha}; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A_{\text{free}}, A_{\delta, \alpha}\}$ admits the representation

$$\begin{aligned} & S(A_{\text{free}}, A_{\delta, \alpha}; \lambda) \\ &= I_{\mathcal{H}_\lambda} + 2i \sqrt{\text{Im} \Lambda_1(\lambda + i0)^{-1}} (I - \alpha \Lambda_1(\lambda + i0)^{-1})^{-1} \alpha \sqrt{\text{Im} \Lambda_1(\lambda + i0)^{-1}}. \end{aligned}$$

Proof. Note first that the assumptions $\alpha^{-1} \in L^\infty(\mathcal{C})$ implies that the closed symmetric operator $S = A_{\text{free}} \cap A_{\delta, \alpha}$ is given by

$$Sf = \mathcal{L}f, \\ \text{dom}(S) = \{f = \{f^+, f^-\} \in H^2(\mathbb{R}^n) : \gamma_D^+ f^+ = \gamma_D^- f^- = 0\}$$

and hence coincides with the one in (5.9) (in the case $n = 2$). It follows from [21, Corollary 4.4] that the operator S is simple and as in the proof of Theorem 5.1 one verifies that its adjoint S^* is given by

$$S^*f = \mathcal{L}f, \\ \text{dom}(S^*) = \{f = \{f^+, f^-\} \in H_\Delta^0(\mathbb{R}^n \setminus \mathcal{C}) : \gamma_D^+ f^+ = \gamma_D^- f^-\}.$$

Next we define the operator T by

$$Tf = \mathcal{L}f, \\ \text{dom}(T) = \{f = \{f^+, f^-\} \in H_\Delta^{3/2}(\mathbb{R}^n \setminus \mathcal{C}) : \gamma_D^+ f^+ = \gamma_D^- f^-\} \quad (5.27)$$

and for $f = \{f^+, f^-\} \in \text{dom}(T)$ we write $\gamma_D f := \gamma_D^+ f^+ = \gamma_D^- f^-$ as in (5.11). We will show that $\Pi_{\delta, \alpha}^{\text{free}} = \{L^2(\mathcal{C}), \Gamma_0, \Gamma_1\}$, where

$$\Gamma_0 f = \gamma_N^+ f^+ + \gamma_N^- f^-, \quad f \in \text{dom}(T),$$

and

$$\Gamma_1 f = \gamma_D f - \frac{1}{\alpha} (\gamma_N^+ f^+ + \gamma_N^- f^-), \quad f \in \text{dom}(T),$$

is a B -generalized boundary triple such that

$$A_{\text{free}} = T \upharpoonright \ker(\Gamma_0) \quad \text{and} \quad A_{\delta, \alpha} = T \upharpoonright \ker(\Gamma_1), \quad (5.28)$$

and the corresponding Weyl function

$$M_{\delta, \alpha}^{\text{free}}(z) := \Lambda_1(z)^{-1} - \frac{1}{\alpha}, \quad z \in \mathbb{C}_+, \quad (5.29)$$

is \mathfrak{S}_1 -regular.

In fact, for $f = \{f^+, f^-\}$, $g = \{g^+, g^-\} \in \text{dom}(T)$ we compute with the help of Green's identity (4.6) and the interface conditions $\gamma_D^+ f^+ = \gamma_D^- f^-$ and $\gamma_D^+ g^+ = \gamma_D^- g^-$ that

$$\begin{aligned}
& (\Gamma_1 f, \Gamma_0 g) - (\Gamma_0 f, \Gamma_1 g) \\
&= (\gamma_D f - \alpha^{-1}(\gamma_N^+ f^+ + \gamma_N^- f^-), \gamma_N^+ g^+ + \gamma_N^- g^-) \\
&\quad - (\gamma_N^+ f^+ + \gamma_N^- f^-, \gamma_D g - \alpha^{-1}(\gamma_N^+ g^+ + \gamma_N^- g^-)) \\
&= (\gamma_D f, \gamma_N^+ g^+ + \gamma_N^- g^-) - (\gamma_N^+ f^+ + \gamma_N^- f^-, \gamma_D g) \\
&= (\gamma_D^+ f^+, \gamma_N^+ g^+) - (\gamma_N^+ f^+, \gamma_D^+ g^+) + (\gamma_D^- f^-, \gamma_N^- g^-) - (\gamma_N^- f^-, \gamma_D^- g^-) \\
&= (-\Delta f^+, g^+) - (f^+, -\Delta g^+) + (-\Delta f^-, g^-) - (f^-, -\Delta g^-) \\
&= (Tf, g) - (f, Tg),
\end{aligned}$$

which shows (2.1). In order to show that Γ_0 is surjective we fix some $\lambda_0 \in \mathbb{R}$ such that $\lambda_0 < \min\{\sigma(A_D), \sigma(A_N)\}$ and note that the direct sum decomposition

$$\operatorname{dom}(T) = \operatorname{dom}(A_D) \dot{+} \ker(T - \lambda_0)$$

holds since $\lambda_0 \in \rho(A_D)$. It follows from (5.27) and (4.4) that γ_D maps $\ker(T - \lambda_0)$ onto $H^1(\mathcal{C})$. As $\Lambda_1^\pm(\lambda_0) = (\mathcal{N}^\pm(\lambda_0))^{-1}$ (cf. (4.29)) are uniformly positive self-adjoint operators in $L^2(\mathcal{C})$, it follows that also $\Lambda_1(\lambda_0) = \Lambda_1^+(\lambda_0) + \Lambda_1^-(\lambda_0)$ is a uniformly positive self-adjoint operator in $L^2(\mathcal{C})$. Let $\psi \in L^2(\mathcal{C})$, choose $\varphi \in H^1(\mathcal{C})$ and $f_{\lambda_0} = \{f_{\lambda_0}^+, f_{\lambda_0}^-\} \in \ker(T - \lambda_0)$ such that $\Lambda_1(\lambda_0)\varphi = \psi$ and $\gamma_D f_{\lambda_0} = \varphi$. Then we have

$$\Gamma_0 f_{\lambda_0} = \gamma_N^+ f_{\lambda_0}^+ + \gamma_N^- f_{\lambda_0}^- = \Lambda_1(\lambda_0) \gamma_D f_{\lambda_0} = \Lambda_1(\lambda_0) \varphi = \psi$$

and this implies $\operatorname{ran}(\Gamma_0) = L^2(\mathcal{C})$.

It is not difficult to check that $\operatorname{dom}(A_{\text{free}})$ and $\operatorname{dom}(A_{\delta, \alpha})$ are contained in $\ker(\Gamma_0)$ and $\ker(\Gamma_1)$, respectively, and since A_{free} and $A_{\delta, \alpha}$ are self-adjoint and $T \upharpoonright \ker(\Gamma_0)$ and $T \upharpoonright \ker(\Gamma_1)$ are symmetric by Green's identity (2.1) it follows that (5.28) holds. Since $S = A_{\text{free}} \cap A_{\delta, \alpha}$ and

$$(\operatorname{dom}(A_{\text{free}}) + \operatorname{dom}(A_{\delta, \alpha})) \subset \operatorname{dom}(T) \subset \operatorname{dom}(S^*),$$

Proposition 2.9 implies $\overline{T} = S^*$. Hence $\Pi_{\delta, \alpha}^{\text{free}}$ is a B -generalized boundary triple such that (5.28) is satisfied.

In order to show that the corresponding Weyl function is given by (5.29) let $f_z = \{f_z^+, f_z^-\} \in \ker(T - z)$ and $z \in \mathbb{C}_+$. Then we have

$$\Lambda_1(z) \gamma_D f_z = \Lambda_1^+(z) \gamma_D^+ f_z^+ + \Lambda_1^-(z) \gamma_D^- f_z^- = \gamma_N^+ f_z^+ + \gamma_N^- f_z^- = \Gamma_0 f_z$$

and since $\ker(\Lambda_1(z)) = \{0\}$ we conclude

$$\left(\Lambda_1(z)^{-1} - \frac{1}{\alpha} \right) \Gamma_0 f_z = \gamma_D f_z - \frac{1}{\alpha} (\gamma_N^+ f_z^+ - \gamma_N^- f_z^-) = \Gamma_1 f_z.$$

This proves the representation (5.29). In order to see that the Weyl function $M_{\delta,\alpha}^{\text{free}}(\cdot)$ is \mathfrak{S}_1 -regular we argue in the same way as in the previous proofs. Denote the γ -field corresponding to the B -generalized boundary triple $\Pi_{\delta,\alpha}^{\text{free}}$ by $\gamma_{\delta,\alpha}^{\text{free}}(\cdot)$ and use

$$M_{\delta,\alpha}^{\text{free}}(z) = M_{\delta,\alpha}^{\text{free}}(\xi)^* + (z - \xi)\gamma_{\delta,\alpha}^{\text{free}}(\xi)^*\gamma_{\delta,\alpha}^{\text{free}}(z) \quad (5.30)$$

with some $\xi \in \rho(A_{\text{free}}) \cap \rho(A_{\delta,\alpha}) \cap \mathbb{R}$ and all $z \in \rho(A_{\text{free}})$. For $h = \{h^+, h^-\} \in L^2(\mathbb{R}^n)$ we have

$$\gamma_{\delta,\alpha}^{\text{free}}(\xi)^*h = \Gamma_1(A_{\text{free}} - \xi)^{-1}h = \gamma_D(A_{\text{free}} - \xi)^{-1}h \in H^{3/2}(\mathcal{C})$$

and hence Lemma 4.1 applied with $r = 3/2$ and $s = 0$ yields

$$\gamma_{\delta,\alpha}^{\text{free}}(\xi)^* \in \mathcal{S}_{\frac{2(n-1)}{3}}(L^2(\mathbb{R}^n), L^2(\mathcal{C})). \quad (5.31)$$

As before we conclude

$$\gamma_{\delta,\alpha}^{\text{free}}(z) \in \mathcal{S}_{\frac{2(n-1)}{3}}(L^2(\mathcal{C}), L^2(\mathbb{R}^n)), \quad z \in \rho(A_{\text{free}}). \quad (5.32)$$

It follows from (5.30) that

$$K(z) := M_{\delta,\alpha}^{\text{free}}(z) - M_{\delta,\alpha}^{\text{free}}(\xi) \in \mathcal{S}_{\frac{n-1}{3}}(L^2(\mathcal{C})) \subset \mathfrak{S}_1(L^2(\mathcal{C})), \quad z \in \mathbb{C}_+,$$

where $M_{\delta,\alpha}^{\text{free}}(\xi) = M_{\delta,\alpha}^{\text{free}}(\xi)^*$ was used. Since the operator $\frac{1}{\alpha}$ is boundedly invertible and $\text{ran}(\Lambda_1(\xi)^{-1}) \subseteq H^1(\mathcal{C})$, the operator $M_{\delta,\alpha}^{\text{free}}(\xi)$ is a Fredholm operator. Furthermore, $\ker(M_{\delta,\alpha}^{\text{free}}(\xi)) = \{0\}$ by Lemma 2.14 for $\xi \in \rho(A_{\text{free}}) \cap \rho(A_{\delta,\alpha}) \cap \mathbb{R}$. Hence $M_{\delta,\alpha}^{\text{free}}(\xi)$ is boundedly invertible and it follows that $M_{\delta,\alpha}^{\text{free}}(\cdot)$ is \mathfrak{S}_1 -regular for $n = 2, 3$.

The assertions in Theorem 5.6 follow from Theorem 3.1 and relations

$$\text{Im } M_{\delta,\alpha}^{\text{free}}(z) = \text{Im } \Lambda_1(z), \quad M_{\delta,\alpha}^{\text{free}}(z)^{-1} = -(I - \alpha\Lambda_1(z)^{-1})^{-1}\alpha, \quad z \in \mathbb{C}_+,$$

and

$$\begin{aligned} \text{Im } M_{\delta,\alpha}^{\text{free}}(\lambda + i0) &= \text{Im } \Lambda_1(\lambda + i0), \\ M_{\delta,\alpha}^{\text{free}}(\lambda + i0)^{-1} &= -(I - \alpha\Lambda_1(\lambda + i0)^{-1})^{-1}\alpha \end{aligned}$$

for a.e. $\lambda \in \mathbb{R}$. \square

Remark 5.7. As in previous remarks it follows from (5.31)–(5.32) and Krein's formula in Proposition 2.7 (iii) that

$$(A_{\delta,\alpha} - z)^{-1} - (A_{\text{free}} - z)^{-1} = -\gamma_{\delta,\alpha}^{\text{free}}(z)M_{\delta,\alpha}^{\text{free}}(z)^{-1}\gamma_{\delta,\alpha}^{\text{free}}(\bar{z})^* \in \mathcal{S}_{\frac{n-1}{3}}(L^2(\mathbb{R}^n))$$

for all $z \in \rho(A_{\text{free}}) \cap \rho(A_{\delta,\alpha})$; cf. [16].

Acknowledgments

The authors are indebted to Ludvig Faddeev, Boris Pavlov, and Andrea Posilicano for useful discussions and remarks. Jussi Behrndt gratefully acknowledges financial support by the Austrian Science Fund (FWF), project P 25162-N26. The preparation of the paper was supported by the European Research Council via ERC-2010-AdG no. 267802 (“Analysis of Multiscale Systems Driven by Functionals”). The publication was financially supported by the Ministry of Education and Science of the Russian Federation (the Agreement number No. 02.A03.21.0008).

Appendix A. Spectral representation and scattering matrix

A.1. Spectral representations and operator spectral integrals

Let $E(\cdot)$ be a spectral measure in the separable Hilbert space \mathfrak{H} defined on the Borel sets $\mathfrak{B}(\mathbb{R})$ of the real axis \mathbb{R} . Further, let C be a Hilbert–Schmidt operator in \mathfrak{H} . Obviously, $\Sigma(\delta) := C^*E(\delta)C$, $\delta \in \mathfrak{B}(\mathbb{R})$ defines a trace class valued measure on $\mathfrak{B}(\mathbb{R})$ of finite variation; cf. [12, Lemma 3.11]. The measure admits a unique decomposition

$$\Sigma(\cdot) = \Sigma^s(\cdot) + \Sigma^{ac}(\cdot)$$

into a singular measure $\Sigma^s(\cdot) = C^*E^s(\cdot)C$ and an absolutely continuous measure $\Sigma^{ac}(\cdot) = C^*E^{ac}(\cdot)C$. From [12, Proposition 3.13] it follows that the trace class valued function $\Sigma(\lambda) := C^*E((-\infty, \lambda))C$ admits a derivative $K(\lambda) := \frac{d}{d\lambda}\Sigma(\lambda) \geq 0$ in the trace class norm for a.e. $\lambda \in \mathbb{R}$ with respect the Lebesgue measure $d\lambda$ such that

$$\Sigma^{ac}(\delta) = \int_{\delta} K(\lambda) d\lambda, \quad \delta \in \mathfrak{B}(\mathbb{R}).$$

By $\mathcal{H}_{\lambda} := \overline{\text{ran}(K(\lambda))} \subseteq \mathfrak{H}$ we define a measurable family of subspaces in \mathfrak{H} . The orthogonal projection $P(\lambda)$ from \mathfrak{H} onto \mathcal{H}_{λ} form a measurable family of projections which defines by

$$(Pf)(\lambda) := P(\lambda)f(\lambda), \quad f \in L^2(\mathbb{R}, d\lambda, \mathfrak{H}),$$

an orthogonal projection from $L^2(\mathbb{R}, d\lambda, \mathfrak{H})$ onto a subspace which is denoted by $L^2(\mathbb{R}, d\lambda, \mathcal{H}_{\lambda})$. Let us assume that the closed linear span of the sets $E^{ac}(\delta)\text{ran}(C)$, $\delta \in \mathfrak{B}(\mathbb{R})$, coincides with $\mathfrak{H}^{ac} = E^{ac}(\mathbb{R})\mathfrak{H}$. Let

$$(\Phi E^{ac}(\delta)Cf)(\lambda) := \chi_{\delta}(\lambda)\sqrt{K(\lambda)}f, \quad \delta \in \mathfrak{B}(\mathbb{R}), \quad f \in \mathfrak{H},$$

where $\chi_{\delta}(\cdot)$ denotes the characteristic function of $\delta \in \mathfrak{B}(\mathbb{R})$. Obviously, we have

$$\int \|(\Phi E^{ac}(\delta)Cf)(\lambda)\|_{\mathfrak{H}}^2 d\lambda = \int_{\delta} \|\sqrt{K(\lambda)}f\|_{\mathfrak{H}}^2 d\lambda = \|E^{ac}(\delta)Cf\|_{\mathfrak{H}}^2.$$

Hence $\Phi : \mathfrak{H}^{ac} \rightarrow L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$ defines an isometry from \mathfrak{H}^{ac} into $L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$. Let us show that Φ is onto $L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$. Let $g \in L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$ such that

$$0 = (\Phi E^{ac}(\delta)Cf, g) = \int_{\delta} (\sqrt{K(\lambda)}f, g(\lambda))_{\mathfrak{H}} d\lambda$$

for $f \in \mathfrak{H}^{ac}$, $\delta \in \mathfrak{B}(\mathbb{R})$. Since δ is arbitrary we find $(\sqrt{K(\lambda)}f, g(\lambda))_{\mathfrak{H}} = 0$ for a.e. $\lambda \in \mathbb{R}$. Hence $g(\lambda) \perp \mathcal{H}_\lambda$ for a.e. $\lambda \in \mathbb{R}$ which shows $g(\lambda) = 0$ for a.e. $\lambda \in \mathbb{R}$. Hence Φ is an isometry from \mathfrak{H}^{ac} onto the subspace $L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$.

Obviously, we have

$$(\Phi E^{ac}(\delta)f)(\lambda) = \chi_\delta(\lambda)(\Phi f)(\lambda), \quad \delta \in \mathfrak{B}(\mathbb{R}), \quad f \in \mathfrak{H}^{ac}.$$

Let A be a self-adjoint operator in \mathfrak{H} and let $E_A(\cdot)$ be the corresponding spectral measure, i.e. $A = \int_{\mathbb{R}} \lambda dE_A(\lambda)$. Then $M\Phi = \Phi A^{ac}$ where M is the natural multiplication operator defined by

$$(Mf)(\lambda) := \lambda f(\lambda),$$

$$f \in \text{dom}(M) := \{f \in L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda) : \lambda f(\lambda) \in L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)\}.$$

If $\varphi(\cdot) : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded Borel function then $\varphi(M)\Phi = \Phi\varphi(A^{ac})$.

Lemma A.1. *Let A , $E_A(\cdot)$, C and $K(\lambda)$ be as above and assume that the absolutely continuous subspace $\mathfrak{H}^{ac}(A)$ satisfies the condition*

$$\mathfrak{H}^{ac}(A) = \text{clsp} \{E_A^{ac}(\delta)\text{ran}(C) : \delta \in \mathfrak{B}(\mathbb{R})\}.$$

Then the mapping

$$E^{ac}(\delta)Cf \mapsto \chi_\delta(\lambda)\sqrt{K(\lambda)}f \quad \text{for a.e. } \lambda \in \mathbb{R}, \quad f \in \mathfrak{H},$$

onto the dense subspace $\text{span}\{E_A^{ac}(\delta)\text{ran}(C) : \delta \in \mathfrak{B}(\mathbb{R})\}$ of $\mathfrak{H}^{ac}(A)$ admits a unique continuation to an isometric isomorphism from $\Phi : \mathfrak{H}^{ac}(A) \rightarrow L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$ such that

$$(\Phi E_A^{ac}(\delta)g)(\lambda) = \chi_\delta(\lambda)(\Phi g)(\lambda), \quad g \in \mathfrak{H}^{ac}(A),$$

holds for any $\delta \in \mathfrak{B}(\mathbb{R})$.

Let us consider operator spectral integrals of the form $\int_{\mathbb{R}} dE^{ac}(\mu)Cf(\lambda)$, which are defined whenever $f(\cdot) : \mathbb{R} \rightarrow \mathfrak{H}$ is a Borel measurable function, cf. [12, Section 5.2]. From

[12, Proposition 5.13] we find that this integral exists if and only if $\int_{\mathbb{R}} \|\sqrt{K(\mu)}f(\mu)\|_{\mathfrak{H}}^2 d\mu$ exists and is finite. One verifies that

$$\left(\Phi \int_{\mathbb{R}} dE^{ac}(\mu) C f(\mu) \right) (\lambda) = \sqrt{K(\lambda)} f(\lambda). \quad (\text{A.1})$$

A.2. Scattering

In the following let A and B be self-adjoint operators in \mathfrak{H} , let $J \in \mathcal{L}(\mathfrak{H})$ be a bounded operator such that $J \operatorname{dom} A \subseteq \operatorname{dom} B$. If

$$V := BJ - JA, \quad \operatorname{dom} V := \operatorname{dom} A,$$

is closable and its closure is a trace class operator then the wave operators

$$W_{\pm}(A, B; J) := s - \lim_{t \rightarrow \pm\infty} e^{itB} J e^{-itA} P^{ac}(A)$$

exist, see [12, 71, 73]. The scattering operator S_J is defined by

$$S_J(A, B) := W_+(A, B; J)^* W_-(A, B; J).$$

Usually the wave operators $W_{\pm}(A, B; J)$ and the scattering operator S_J are not the quantities of main interest. The objects one is more interested in are the wave operators $W_{\pm}(A, B) := W_{\pm}(A, B; I)$ and $S(A, B) := S_I(A, B)$. However, if the resolvent difference of A and B is trace class, then the existence of $W_{\pm}(A, B; J)$ with $J = -(B - i)^{-1}(A - i)^{-1}$ yields the existence of $W_{\pm}(A, B)$ and both operators are related by

$$W_{\pm}(A, B; J) = -W_{\pm}(A, B)(A - i)^{-2}.$$

In particular, this yields

$$S_J(A, B) = S(A, B)(I + A^2)^{-2}. \quad (\text{A.2})$$

The following theorem was announced in [20, Appendix A] but not proved there. Below the complete proof of this theorem is given.

Theorem A.2. *Let A and B be self-adjoint operators in the separable Hilbert space \mathfrak{H} and suppose that the resolvent difference admits the factorization*

$$\mathfrak{S}_1(\mathfrak{H}) \ni (B - i)^{-1} - (A - i)^{-1} = \phi(A)CGC^* = QC^*, \quad (\text{A.3})$$

where $C \in \mathfrak{S}_2(\mathcal{H}, \mathfrak{H})$, $G \in \mathcal{L}(\mathcal{H})$, $\phi(\cdot) : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded continuous function and $Q = \phi(A)CG$. Assume that the condition

$$\mathfrak{H}^{ac}(A) = \text{clsp} \{E_A^{ac}(\delta) \text{ran}(C) : \delta \in \mathfrak{B}(\mathbb{R})\} \quad (\text{A.4})$$

is satisfied and let $K(\lambda) = \frac{d}{d\lambda} C^* E_A((-\infty, \lambda)) C$ and $\mathcal{H}_\lambda = \overline{\text{ran}(K(\lambda))}$ for a.e. $\lambda \in \mathbb{R}$. Then $L^2(\mathbb{R}, d\lambda, \mathcal{H}_\lambda)$ is a spectral representation of A^{ac} and the scattering matrix $\{S(A, B; \lambda)\}_{\lambda \in \mathbb{R}}$ of the scattering system $\{A, B\}$ has the representation

$$S(A, B; \lambda) = I_{\mathcal{H}_\lambda} + 2\pi i(1 + \lambda^2)^2 \sqrt{K(\lambda)} Z(\lambda) \sqrt{K(\lambda)} \quad (\text{A.5})$$

for a.e. $\lambda \in \mathbb{R}$, where

$$Z(\lambda) = \frac{1}{\lambda + i} Q^* Q + \frac{\phi(\lambda)}{(\lambda + i)^2} G + \lim_{\varepsilon \rightarrow +0} Q^* (B - (\lambda + i\varepsilon))^{-1} Q \quad (\text{A.6})$$

and the limit of the last term on the right hand side exists in the Hilbert–Schmidt norm.

Proof. Consider the scattering operator

$$S_J(A, B) := W_+(A, B; J)^* W_-(A, B; J) : \mathfrak{H}^{ac}(A) \longrightarrow \mathfrak{H}^{ac}(A),$$

where $J := -R_B(i)R_A(i)$ and

$$R_B(\xi) := (B - \xi)^{-1}, \quad R_A(\xi) := (A - \xi)^{-1}.$$

One easily checks that

$$V := BJ - JA = (B - i)^{-1} - (A - i)^{-1} = \phi(A) C G C^*$$

where we have used the assumption (A.3). We note that the scattering operator commutes with A . From [12, Theorem 18.4] one gets the representation

$$S_J(A, B) - W_+(A, B; J)^* W_+(A, B; J) = s - \lim_{\epsilon \rightarrow +0} w - \lim_{\tau \rightarrow +0} \left\{ -2\pi i \int_{\mathbb{R}} dE_A^{ac}(\lambda) T(\tau; \lambda) \delta_\epsilon(A; \lambda) P^{ac}(A) \right\},$$

where

$$T(\tau; \lambda) := J^* V - V^* R_B(\lambda + i\tau) V$$

and

$$\delta_\epsilon(A; \lambda) := \frac{1}{2\pi i} (R_A(\lambda + i\epsilon) - R_A(\lambda - i\epsilon)) = \frac{1}{\pi} \frac{\epsilon}{(A - \lambda)^2 + \epsilon^2}.$$

If condition (A.3) is satisfied, then

$$R_B(i) = R_A(i) + \phi(A)CGC^* = R_A(i) + QC^*$$

and we get

$$\begin{aligned} J^*V &= -R_A(-i)R_B(-i)V \\ &= -R_A(-i)CQ^*V - R_A(-i)^2V \\ &= -R_A(-i)CQ^*V - R_A(-i)^2\phi(A)CGC^*. \end{aligned}$$

Hence we find

$$T(\tau; \lambda) = -\left(R_A(-i)CQ^*Q + R_A(-i)^2\phi(A)CG + CQ^*R_B(\lambda + i\tau)Q\right)C^*.$$

Using (A.1) we get

$$\begin{aligned} \left(\Phi \int_{\mathbb{R}} dE_A^{ac}(\mu)T(\tau; \mu)\delta_\epsilon(A; \mu)P^{ac}(A)Ch\right)(\lambda) = \\ -\sqrt{K(\lambda)}Z(\tau; \lambda)C^*\delta_\epsilon(A; \lambda)P^{ac}(A)Ch, \end{aligned}$$

where

$$Z(\tau; \lambda) := \frac{1}{\lambda + i}Q^*Q + \frac{\phi(\lambda)}{(\lambda + i)^2}G + Q^*R_B(\lambda + i\tau)Q.$$

We note that the limit $Q^*R_B(\lambda + i0)Q := \lim_{\tau \rightarrow +0} Q^*R_B(\lambda + i\tau)Q$ exists in the Hilbert–Schmidt norm. Hence the limit $Z(\lambda) := \lim_{\tau \rightarrow +0} Z(\tau; \lambda)$ exists in the operator norm and is given by

$$Z(\lambda) = \frac{1}{\lambda + i}Q^*Q + \frac{\phi(\lambda)}{(\lambda + i)^2}G + Q^*R_B(\lambda + i0)Q.$$

This gives

$$\left(\Phi \left\{ \text{s-lim}_{\epsilon \rightarrow +0} \text{w-lim}_{\tau \rightarrow +0} \int_{\mathbb{R}} dE_A^{ac}(\mu)T(\tau; \mu)\delta_\epsilon(A; \mu)P^{ac}(A)Ch \right\}\right)(\lambda) = -\sqrt{K(\lambda)}Z(\lambda)K(\lambda)h.$$

By the compactness of V we get that $W_+(A, B; J)^*W_+(A, B; J) = (I + A^2)^{-2}$. Therefore we have

$$(\Phi(W_+(A, B; J)^*W_+(A, B; J)\Phi^*f)(\lambda) = (1 + \lambda^2)^{-2}f(\lambda).$$

Hence $\Phi S_J(A, B)\Phi^*$ is equal to a multiplication operator with a measurable function $S_J(A, B; \lambda) : \mathcal{H}_\lambda \rightarrow \mathcal{H}_\lambda$ given by

$$S_J(A, B; \lambda) := (1 + \lambda^2)^{-2} I_{\mathcal{H}_\lambda} + 2\pi i \sqrt{K(\lambda)} Z(\lambda) \sqrt{K(\lambda)}.$$

Using (A.2) we find that $\Phi S(A, B) \Phi^*$ is a multiplication operator induced by the measurable function $S(A, B; \lambda) : \mathcal{H}_\lambda \rightarrow \mathcal{H}_\lambda$. Both functions $S_J(A, B; \lambda)$ and $S(A, B; \lambda)$ are related by

$$S_J(A, B; \lambda) = S(A, B; \lambda)(1 + \lambda^2)^{-2}$$

which yields the representation (A.5). \square

References

- [1] H. Abels, G. Grubb, I.G. Wood, Extension theory and Kreĭn-type resolvent formulas for nonsmooth boundary value problems, *J. Funct. Anal.* 266 (2014) 4037–4100.
- [2] V.M. Adamyan, B.S. Pavlov, Zero-radius potentials and M. G. Krein’s formula for generalized resolvents, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 149 (1986) 7–23, translation in *J. Sov. Math.* 42 (1988) 1537–1550.
- [3] S. Agmon, *Lectures on Elliptic Boundary Value Problems*, D. Van Nostrand Co., Princeton, NJ–Toronto–London, 1965.
- [4] M.S. Agranovich, Elliptic operators on closed manifolds, in: *Partial Differential Equations, VI*, in: *Encyclopaedia Math. Sci.*, vol. 63, Springer, Berlin, 1990, pp. 1–130.
- [5] M.S. Agranovich, Elliptic boundary problems, in: *Partial Differential Equations, IX*, in: *Encyclopaedia Math. Sci.*, vol. 79, Springer, Berlin, 1997, pp. 1–144, 275–281.
- [6] A. Albeverio, F. Gesztesy, R. Høegh-Krohn, H. Holden, *Solvable Models in Quantum Mechanics*, second edition, AMS Chelsea Publishing, Providence, RI, 2005, with an appendix by Pavel Exner.
- [7] S. Albeverio, A.S. Kostenko, M.M. Malamud, H. Neidhardt, Spherical Schrödinger operators with δ -type interactions, *J. Math. Phys.* 54 (2013) 052103, 24 pp.
- [8] S. Albeverio, P. Kurasov, *Singular Perturbations of Differential Operators: Solvable Schrödinger Type Operators*, London Mathematical Society Lecture Note Series, vol. 271, Cambridge University Press, Cambridge, 2000.
- [9] W.O. Amrein, D.B. Pearson, M operators: a generalisation of Weyl–Titchmarsh theory, *J. Comput. Appl. Math.* 171 (2004) 1–26.
- [10] J.-P. Antoine, F. Gesztesy, J. Shabani, Exactly solvable models of sphere interactions in quantum mechanics, *J. Phys. A* 20 (1987) 3687–3712.
- [11] N.T. Bagraev, A.B. Mikhailova, B.S. Pavlov, L.V. Prokhorov, A.M. Yafyasov, Parameter regime of a resonance quantum switch, *Phys. Rev. B* 71 (2005) 165308.
- [12] H. Baumgärtel, M. Wollenberg, *Mathematical Scattering Theory, Operator Theory: Advances and Applications*, vol. 9, Birkhäuser, Basel, 1983.
- [13] J. Behrndt, M. Langer, Boundary value problems for elliptic partial differential operators on bounded domains, *J. Funct. Anal.* 243 (2007) 536–565.
- [14] J. Behrndt, M. Langer, Elliptic operators, Dirichlet-to-Neumann maps and quasi boundary triples, in: *Operator Methods for Boundary Value Problems*, in: *London Math. Soc. Lecture Note Ser.*, vol. 404, Cambridge Univ. Press, Cambridge, 2012, pp. 121–160.
- [15] J. Behrndt, M. Langer, I. Lobanov, V. Lotoreichik, I.Yu. Popov, A remark on Schatten–von Neumann properties of resolvent differences of generalized Robin Laplacians on bounded domains, *J. Math. Anal. Appl.* 371 (2010) 750–758.
- [16] J. Behrndt, M. Langer, V. Lotoreichik, Schrödinger operators with δ and δ' -potentials supported on hypersurfaces, *Ann. Henri Poincaré* 14 (2013) 385–423.
- [17] J. Behrndt, M. Langer, V. Lotoreichik, Spectral estimates for resolvent differences of self-adjoint elliptic operators, *Integral Equations Operator Theory* 77 (2013) 1–37.
- [18] J. Behrndt, M. Langer, V. Lotoreichik, J. Rohleder, Quasi boundary triples and semibounded self-adjoint extensions, *Proc. Roy. Soc. Edinburgh Sect. A* (2017), in press.
- [19] J. Behrndt, M.M. Malamud, H. Neidhardt, Scattering matrices and Weyl functions, *Proc. Lond. Math. Soc.* 97 (2008) 568–598.

- [20] J. Behrndt, M.M. Malamud, H. Neidhardt, Finite rank perturbations, scattering matrices and inverse problems, in: *Recent Advances in Operator Theory in Hilbert and Krein Spaces*, in: *Operator Theory: Advances and Applications*, vol. 198, Birkhäuser Verlag, Basel, 2010, pp. 61–85.
- [21] J. Behrndt, J. Rohleder, Spectral analysis of self-adjoint elliptic differential operators, Dirichlet-to-Neumann maps, and abstract Weyl functions, *Adv. Math.* 285 (2015) 1301–1338.
- [22] J. Behrndt, J. Rohleder, Titchmarsh–Weyl theory for Schrödinger operators on unbounded domains, *J. Spectr. Theory* 6 (2016) 67–87.
- [23] Yu.M. Berezanskiĭ, *Selfadjoint Operators in Spaces of Functions of Infinitely Many Variables*, *Translations of Mathematical Monographs*, vol. 63, American Mathematical Society, Providence, RI, 1986.
- [24] M.Š. Birman, Perturbations of the continuous spectrum of a singular elliptic operator by varying the boundary and the boundary conditions, *Vestnik Leningrad. Univ.* 17 (1962) 22–55.
- [25] M.Š. Birman, S.B. Ėntina, Stationary approach in abstract scattering theory, *Izv. Ross. Akad. Nauk Ser. Mat.* 31 (1967) 401–430.
- [26] L. de Branges, Perturbations of self-adjoint transformations, *Amer. J. Math.* 84 (1962) 543–560.
- [27] J.F. Brasche, P. Exner, Yu.A. Kuperin, P. Šeba, Schrödinger operators with singular interactions, *J. Math. Anal. Appl.* 184 (1994) 112–139.
- [28] J.F. Brasche, M.M. Malamud, H. Neidhardt, Weyl function and spectral properties of self-adjoint extensions, *Integral Equations Operator Theory* 43 (2002) 264–289.
- [29] B.M. Brown, G. Grubb, I.G. Wood, M -functions for closed extensions of adjoint pairs of operators with applications to elliptic boundary problems, *Math. Nachr.* 282 (2009) 314–347.
- [30] B.M. Brown, M. Marletta, S.N. Naboko, I.G. Wood, Boundary triplets and M -functions for non-self-adjoint operators, with applications to elliptic PDEs and block operator matrices, *J. Lond. Math. Soc.* 77 (2008) 700–718.
- [31] J. Brüning, V. Geyler, K. Pankrashkin, Spectra of self-adjoint extensions and applications to solvable Schrödinger operators, *Rev. Math. Phys.* 20 (2008) 1–70.
- [32] J.W. Calkin, Abstract symmetric boundary conditions, *Trans. Amer. Math. Soc.* 45 (1939) 369–442.
- [33] P. Deift, B. Simon, On the decoupling of finite singularities from the question of asymptotic completeness in two body quantum systems, *J. Funct. Anal.* 23 (1976) 218–238.
- [34] V.A. Derkach, S. Hassi, M.M. Malamud, H.S.V. de Snoo, Boundary relations and their Weyl families, *Trans. Amer. Math. Soc.* 358 (2006) 5351–5400.
- [35] V.A. Derkach, S. Hassi, M.M. Malamud, H.S.V. de Snoo, Boundary triplets and Weyl functions. Recent developments, in: *Operator Methods for Boundary Value Problems*, in: *London Math. Soc. Lecture Note Ser.*, vol. 404, Cambridge Univ. Press, Cambridge, 2012, pp. 161–220.
- [36] V.A. Derkach, M.M. Malamud, Weyl function of a Hermitian operator and its connection with characteristic function, preprint (1985), see also, arXiv:1503.08956.
- [37] V.A. Derkach, M.M. Malamud, Generalized resolvents and the boundary value problems for Hermitian operators with gaps, *J. Funct. Anal.* 95 (1991) 1–95.
- [38] V.A. Derkach, M.M. Malamud, The extension theory of Hermitian operators and the moment problem, *J. Math. Sci.* 73 (1995) 141–242.
- [39] P. Exner, Leaky quantum graphs: a review, in: *Analysis on Graphs and its Applications*, in: *Proc. Sympos. Pure Math.*, vol. 77, Amer. Math. Soc., Providence, RI, 2008, pp. 523–564.
- [40] P. Exner, T. Ichinose, Geometrically induced spectrum in curved leaky wires, *J. Phys. A* 34 (2001) 1439–1450.
- [41] P. Exner, S. Kondej, Bound states due to a strong δ interaction supported by a curved surface, *J. Phys. A* 36 (2003) 443–457.
- [42] P. Exner, S. Kondej, Scattering by local deformations of a straight leaky wire, *J. Phys. A* 38 (2005) 4865–4874.
- [43] P. Exner, K. Yoshitomi, Asymptotics of eigenvalues of the Schrödinger operator with a strong δ -interaction on a loop, *J. Geom. Phys.* 41 (2002) 344–358.
- [44] F. Gesztesy, K.A. Makarov, S.N. Naboko, The spectral shift operator, *Operator Theory: Advances and Applications* 108 (1999) 59–90.
- [45] F. Gesztesy, M.M. Malamud, Spectral theory of elliptic operators in exterior domains, arXiv: 0810.1789.
- [46] F. Gesztesy, M. Mitrea, Generalized Robin boundary conditions, Robin-to-Dirichlet maps, and Krein-type resolvent formulas for Schrödinger operators on bounded Lipschitz domains, in: *Perspectives in Partial Differential Equations, Harmonic Analysis and Applications*, in: *Proc. Sympos. Pure Math.*, vol. 79, Amer. Math. Soc., Providence, RI, 2008, pp. 105–173.
- [47] F. Gesztesy, M. Mitrea, Nonlocal Robin Laplacians and some remarks on a paper by Filonov on eigenvalue inequalities, *J. Differential Equations* 247 (2009) 2871–2896.

- [48] F. Gesztesy, M. Mitrea, Robin-to-Robin maps and Krein-type resolvent formulas for Schrödinger operators on bounded Lipschitz domains, *Operator Theory: Advances and Applications* 191 (2009) 81–113.
- [49] F. Gesztesy, M. Mitrea, A description of all self-adjoint extensions of the Laplacian and Krein-type resolvent formulas on non-smooth domains, *J. Anal. Math.* 113 (2011) 53–172.
- [50] F. Gesztesy, M. Mitrea, M. Zinchenko, Variations on a theme of Jost and Pais, *J. Funct. Anal.* 253 (2007) 399–448.
- [51] V.I. Gorbachuk, M.L. Gorbachuk, *Boundary Value Problems for Operator Differential Equations, Mathematics and its Applications: Soviet Series*, vol. 48, Kluwer Academic Publishers Group, Dordrecht, 1991.
- [52] G. Grubb, A characterization of the non-local boundary value problems associated with an elliptic operator, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 22 (1968) 425–513.
- [53] G. Grubb, Properties of normal boundary problems for elliptic even-order systems, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 1 (1974) 1–61.
- [54] G. Grubb, Singular Green operators and their spectral asymptotics, *Duke Math. J.* 51 (1984) 477–528.
- [55] G. Grubb, Krein resolvent formulas for elliptic boundary problems in nonsmooth domains, *Rend. Semin. Mat. Univ. Politec. Torino* 66 (2008) 271–297.
- [56] G. Grubb, The mixed boundary value problem, Krein resolvent formulas and spectral asymptotic estimates, *J. Math. Anal. Appl.* 382 (2011) 339–363.
- [57] G. Grubb, Perturbation of essential spectra of exterior elliptic problems, *Appl. Anal.* 90 (2011) 103–123.
- [58] G. Grubb, Spectral asymptotics for Robin problems with a discontinuous coefficient, *J. Spectr. Theory* 1 (2011) 155–177.
- [59] T. Kato, *Perturbation Theory for Linear Operators*, Springer, Berlin, 1976.
- [60] M.G. Krein, Basic propositions of the theory of representation of Hermitian operators with deficiency index (m, m) , *Ukrainian Math. J.* 1 (1949) 3–66.
- [61] O.A. Ladyzhenskaya, N.N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, New York–London, 1968.
- [62] J.-L. Lions, E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications*, vol. I, Springer, New York, 1972.
- [63] M.M. Malamud, Spectral theory of elliptic operators in exterior domains, *Russ. J. Math. Phys.* 17 (2010) 96–125.
- [64] M.M. Malamud, H. Neidhardt, Perturbation determinants for singular perturbations, *Russ. J. Math. Phys.* 21 (2014) 55–98.
- [65] A. Mantile, A. Posilicano, M. Sini, Self-adjoint elliptic operators with boundary conditions on not closed hypersurfaces, *J. Differential Equations* 261 (2016) 1–55.
- [66] A. Mantile, A. Posilicano, M. Sini, Limiting absorption principle, generalized eigenfunction expansions and scattering matrix for Laplace operators with boundary conditions on hypersurfaces, *J. Spectr. Theory*, in press.
- [67] M. Marletta, Eigenvalue problems on exterior domains and Dirichlet to Neumann maps, *J. Comput. Appl. Math.* 171 (2004) 367–391.
- [68] W. McLean, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, 2000.
- [69] A.B. Mikhailova, B.S. Pavlov, L.V. Prokhorov, Intermediate Hamiltonian via Glazman's splitting and analytic perturbation for meromorphic matrix-functions, *Math. Nachr.* 280 (2007) 1376–1416.
- [70] S.N. Naboko, On the boundary values of analytic operator-valued functions with a positive imaginary part, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 157 (1987) 55–69.
- [71] H. Neidhardt, Zwei-Raum-Verallgemeinerung des Theorems von Rosenblum und Kato, *Math. Nachr.* 84 (1978) 195–211.
- [72] B.S. Pavlov, I. Antoniou, Jump-start in the analytic perturbation procedure for the Friedrichs model, *J. Phys. A* 38 (2005) 4811–4823.
- [73] D.B. Pearson, A generalization of the Birman trace theorem, *J. Funct. Anal.* 28 (1978) 182–186.
- [74] A. Posilicano, Self-adjoint extensions of restrictions, *Oper. Matrices* 2 (2008) 483–506.
- [75] A. Posilicano, Markovian extensions of symmetric second order elliptic differential operators, *Math. Nachr.* 287 (2014) 1848–1885.
- [76] A. Posilicano, L. Raimondi, Krein's resolvent formula for self-adjoint extensions of symmetric second-order elliptic differential operators, *J. Phys. A* 42 (2009) 015204, 11 pages.
- [77] O. Post, Boundary pairs associated with quadratic forms, *Math. Nachr.* 289 (2016) 1052–1099.

- [78] M. Reed, B. Simon, *Methods of Modern Mathematical Physics I. Functional Analysis*, Academic Press, New York–London, 1972.
- [79] M. Reed, B. Simon, *Methods of Modern Mathematical Physics III. Scattering Theory*, Academic Press, New York–London, 1979.
- [80] K. Schmüdgen, *Unbounded Self-adjoint Operators on Hilbert Space*, Graduate Texts in Mathematics, vol. 265, Springer, Dordrecht, 2012.
- [81] J. Weidmann, *Lineare Operatoren in Hilberträumen, Teil II*, B.G. Teubner, Stuttgart, 2003.
- [82] D.R. Yafaev, *Mathematical Scattering Theory*, Translations of Mathematical Monographs, vol. 105, American Mathematical Society, Providence, RI, 1992.